PLANT DISEASE REPORTER INDEX

Issued by

PLANT SCIENCE RESEARCH DIVISION

Plant Industry Station
Volume 54

Beltsville, Maryland
Number 13

AUTHOR INDEX

ABIVARDI, CYRUS, 339, 1085
ACEDO, GREGORIA N., 296
ADAMS, P. B., (114)
ADAMS, ROBERT E., 923
ADENIJI, M. O., 23, 547
AFFELTRANGER, C. E., (560)
AGRIOS, G. N., (296)
AGUIERO, V. M., (565)
AHMAD, SAEED, (630)
AHRENS, R. W., (281)
AL-BELDAWI, A. S., 76, 524
ALDRICH, JANE, 446
ALLEN, D. J., 809
ANDERSON, GERALD W., (460)
ANDREWS, STUART R., (266)
ARIF, ANWAR, 346
ARMEDILLA, A. L., (1007)
ARNY, D. C., 281
ARTMAN, J. D., 799
AULAKH, KIRPAL S., 337
AWODERU, VICTOR A., 520, (1045)
AYERS, J. E., 277, (1123)

BAGGA, H. S., 719, 796
BAGGETT, J. R., (355)
BAILEY, R. L., (19)
BAINES, R. C., (717)
BAKER, RALPH, (446)
BALLANTYNE, BARBARA, 903
BANIHASHEMI, ZIAEDDIN, 261
BAR-JOSEPH, M., 643
BARKSDALE, T. H., 32
BARTELS, R., (851)
BEALS, HAROLD O., (66)
BEARD, JAMES B., (1075)
BECKETT, J. B., (708)
BEHAR, D. S., (1049)
BÉKÉSI, P., 286
BELL, A. A., (408)
BENDA, G. T. A., 815
BEN-JOSEPH, J., (768)
BERAHA, L., 422
BERNARD, R. L., (544)
BESEMER, S., (689)
BEUTE, M. K., 122, 125, (846)
BHARGAVA, K. S., (181), 727
BIEHN, WILLIAM L., 12, 171, 254
BIVENS, J., (187)
BLAZQUEZ, C. H., 52
BLODGETT, EARLE C., 63
BOESEWINKEL, H. J., 504
BOOTH, J. A., 741
BORECKI, Z., 640
BOWEN, H. H., 682
BOYCE, B. R., 638
BOYLE, J. S., (141)
BRADFUTE, O. E., (156)
BRAVERMAN, SAMUEL W., 580
BRENNAN, EILEEN, 704
BRIDGMON, G. H., 547
BROMFIELD, K. R., 381
BROOKS, H. J., (835)
BROWN, DONALD H., 441, 462
BROWN, F. H., 508
BROWNING, J. A., (807)
BUMBIERIS, M., 622
BUTLER, LARRY D., 467
BUTTERFIELD, J. E., (263)
BYNUM, H. H., (488)

CALVERT, O. H., (286)
CAMERON, H. R., 69
CAMPBELL, F. J., (328)
CAMPBELL, FOY, 1118
CAMPBELL, W. A., (419)
CANERDAY, JAMES V., 424
CARLEY, H. EDWIN, 943
CARROLL, R. B., 811
CARTER, ANNE L., (150)
CASPER, R., 851
CEPONIS, M. J., 263, 856, 964
CEVARIO, S. J., (381)
CHAMBERLAIN, J. D., (132)
CHEE, K. H., 897
CHENEY, PHILIP W., 44, 651
CHENG, YENG-HSIUNG, (29)
CHINNADURAI, G., 844
CHRISTIE, R. G., (161)
CHRISTIE, S. R., (161)
CIMANOWSKI, JAN, 81
CLARK, R. L., 624
CLAYTON, J., (635)
COHN, RUTH, (804)
COLE, H., 141, 146, 981, 1080, 1123
COLE, JOHN R., (12)
COLEMAN, OTTO H., (723)
CONVERSE, RICHARD H., 479, 701, 969
COOK, A. A., 893
CORBIN, J. B., (504)
CORDELL, C. E., 560
CORTÉS-MONLLOR, AMELIA, (212)
COX, R. S., 678, 679, 759, 760
COYIER, DUANE L., 647, (651)
COYNE, D. P., 557

PLANT DISEASE REPORTER INDEX

Issued by

PLANT SCIENCE RESEARCH DIVISION

Plant Industry Station
Volume 54

Beltsville, Maryland
Number 13

AUTHOR INDEX

ABIVARDI, CYRUS, 339, 1085
ACEDO, GREGORIA N., 296
ADAMS, P. B., (114)
ADAMS, ROBERT E., 923
ADENIJI, M. O., 23, 547
AFFELTRANGER, C. E., (560)
AGRIOS, G. N., (296)
AGUIERO, V. M., (565)
AHMAD, SAEED, (630)
AHRENS, R. W., (281)
AL-BELDAWI, A. S., 76, 524
ALDRICH, JANE, 446
ALLEN, D. J., 809
ANDERSON, GERALD W., (460)
ANDREWS, STUART R., (266)
ARIF, ANWAR, 346
ARMEDILLA, A. L., (1007)
ARNY, D. C., 281
ARTMAN, J. D., 799
AULAKH, KIRPAL S., 337
AWODERU, VICTOR A., 520, (1045)
AYERS, J. E., 277, (1123)

BAGGA, H. S., 719, 796
BAGGETT, J. R., (355)
BAILEY, R. L., (19)
BAINES, R. C., (717)
BAKER, RALPH, (446)
BALLANTYNE, BARBARA, 903
BANIHASHEMI, ZIAEDDIN, 261
BAR-JOSEPH, M., 643
BARKSDALE, T. H., 32
BARTELS, R., (851)
BEALS, HAROLD O., (66)
BEARD, JAMES B., (1075)
BECKETT, J. B., (708)
BEHAR, D. S., (1049)
BÉKÉSI, P., 286
BELL, A. A., (408)
BENDA, G. T. A., 815
BEN-JOSEPH, J., (768)
BERAHA, L., 422
BERNARD, R. L., (544)
BESEMER, S., (689)
BEUTE, M. K., 122, 125, (846)
BHARGAVA, K. S., (181), 727
BIEHN, WILLIAM L., 12, 171, 254
BIVENS, J., (187)
BLAZQUEZ, C. H., 52
BLODGETT, EARLE C., 63
BOESEWINKEL, H. J., 504
BOOTH, J. A., 741
BORECKI, Z., 640
BOWEN, H. H., 682
BOYCE, B. R., 638
BOYLE, J. S., (141)
BRADFUTE, O. E., (156)
BRAVERMAN, SAMUEL W., 580
BRENNAN, EILEEN, 704
BRIDGMON, G. H., 547
BROMFIELD, K. R., 381
BROOKS, H. J., (835)
BROWN, DONALD H., 441, 462
BROWN, F. H., 508
BROWNING, J. A., (807)
BUMBIERIS, M., 622
BUTLER, LARRY D., 467
BUTTERFIELD, J. E., (263)
BYNUM, H. H., (488)

CALVERT, O. H., (286)
CAMERON, H. R., 69
CAMPBELL, F. J., (328)
CAMPBELL, FOY, 1118
CAMPBELL, W. A., (419)
CANERDAY, JAMES V., 424
CARLEY, H. EDWIN, 943
CARROLL, R. B., 811
CARTER, ANNE L., (150)
CASPER, R., 851
CEPONIS, M. J., 263, 856, 964
CEVARIO, S. J., (381)
CHAMBERLAIN, J. D., (132)
CHEE, K. H., 897
CHENEY, PHILIP W., 44, 651
CHENG, YENG-HSIUNG, (29)
CHINNADURAI, G., 844
CHRISTIE, R. G., (161)
CHRISTIE, S. R., (161)
CIMANOWSKI, JAN, 81
CLARK, R. L., 624
CLAYTON, J., (635)
COHN, RUTH, (804)
COLE, H., 141, 146, 981, 1080, 1123
COLE, JOHN R., (12)
COLEMAN, OTTO H., (723)
CONVERSE, RICHARD H., 479, 701, 969
COOK, A. A., 893
CORBIN, J. B., (504)
CORDELL, C. E., 560
CORTÉS-MONLLOR, AMELIA, (212)
COX, R. S., 678, 679, 759, 760
COYIER, DUANE L., 647, (651)
COYNE, D. P., 557

CRANDALL, P. C., 132
CRANE, J. L., (1095)
CRAWFORD, JOHNNY L., 324, (586)
CROSIER, D. C., (349), (783), (927)
CROSIER, W. F., 349, 783, 927
CUTRIGHT, NOEL J., 771, 1018
CZABATOR, F. J., 19, 498

DAINES, ROBERT H., 486, (704), 764
DAVIS, D. J., (749)
DAVIS, D. R., 474
DAVIS, DONALD D., 929, 931
DAVIS, S. H., Jr., (606), (1037)
DAVIS, TERRY C., 66
DEMSKI, JAMES W., 880
DENNIS, R. E., (1064)
DESAI, B. G., 976
DICKENS, LESTER E., 266
DIMOND, A. E., (12)
DISHON, IRENE, (232), (909)
DJAKAMIHARDJA, SJAMSUDIN, 307
DOCAMPO, DELIA M., 939
DRAKE, CHARLES R., 686
DRAPER, A. D., (362), 739
DRIVER, CHAS. H., 330
DUCLOS, LEO A., (319), (544)
DUICH, J., (146), (981), (1080)
DUNLEAVY, J. M., (301), 901
DUNN, T. G., (165)

EBRON, T. T., (1045)
ECHANDI, EDDIE, 183
EDMONDS, R. L., (330)
EDWARDSON, J. R., 161
EGLITIS, MAKSIS, 268
ELLINGBOE, A. H., (635)
EL NUR, E., 405, (528)
EMGE, ROBERT G., 130, (512)
EPPS, JAMES M., 319, (544)
EPSTEIN, ABRAHAM H., 679
ESSER, R. P., (692)

FAGBENLE, HELEN, (191)
FARRAR, LUTHER L., 1099
FELDMAN, J. M., 722
FELIX, LEONARD S., 15
FENTON, ROBERT, (869)
FENWICK, H. S., (467)
FERGUS, C. L., (217)
FERRETTI, R. J., 408
FILER, T. H., Jr., 320, (322)
FINNIS, T. M., (330)
FIRMAN, I. D., 1035
FISHER, FRAN. E., 173
FORD, R. E., 191
FORSBERG, J. L., 289, 1095
FOSTER, JANICE G., (251)
FOX, J. A., (544)
FREDERIKSEN, R. A., (311)
FREEMAN, G. C., (952)
FREEMAN, T. E., 358
FREIGOUN S., (405)
FRENCH, D. W., 706
FRENCH, E. R., (315)
FRIDLUND, P. R., 917, 1073
FROSHEISER, F. I., 591
FROWD, J. A., (370), 734
FUCHS, EVA, (347)
FUCIKOVSKY, L. A., 762
FULTON, ROBERT W., 949
FUTRELL, M. C., (307), 311, (483), 1111

GAIROLA, C., 832
GANGOPADHYAY, S., 1088
GARCÍA, JULIO, 403
GARDNER, M. W., 399
GARNSEY, S. M., 907
GARREN, KENNETH H., 840
GAVALAS, N. A., 73
GAY, J. DAN, 243, 604
GECKELER, W. R., (8)
GIANNOPOLITIS, C. N., (561)
GILL, D. L., 1077
GOHEEN, A. C., (1055)
GOLDBERG, C. W., 981, 1080
GOLDEN, A. M., 544
GOLDWEBER, SEYMOUR, (569)
GOODE, M. J., (346)
GOODING, G. V., Jr., 119, (125)
GOODMAN, R. N., (203), (576)
GORDON, D. T., (156)
GORING, C. A. I., (196)
GOULD, CHARLES J., 235, 377
GOVINDASWAMY, C. V., (844)
GOWDA, S. S., 576
GOYAL, J. P., (976)
GRACIA, OLGA, (722)
GRAVES, C. H., Jr., (829)
GREEN, R. J., Jr., 583
GREGORY, GAROLD F., 844
GUDAUSKAS, ROBERT T., (424)
GUMPF, DAVID J., (550)
GUPTA, B. M., 453, 989
GUPTA, O. P., (630), (1092)
GUTTER, Y., 325

HALFON-MEIRI, ALIZA, 442
HALLIWELL, ROBERT S., (669)
HAMPTON, R. O., 355
HANKIN, LESTER, (863)
HARPER, F. C., (368)
HARRIS, H. B., 60
HARRISON, M. B., 306, (771), (1018)
HART, JOHN H., 985
HART, W. H., (1012), (1014), 1016
HARTWIG, E. E., (901)
HAWKSWORTH, FRANK G., (275), 488
HEAGLE, A. S., (8)
HEARN, C. JACK, 869
HEBERT, T. T., (183)
HEGDE, R. K., (788)
HELTON, A. W., 996

HEMPHILL, DONALD V., (488)
HENDRIX, F. F., Jr., 419
HENDRIX, J. WALTER, 347, (384), 387
HEROLD, F., 344
HICKEY, K. D., (395), (792)
HICKMAN, CLINTON E., (923)
HINDS, THOMAS E., 460
HINE, R. B., 179, 1064
HOCKETT, E. A., 749
HODGES, CLINTON F., 206
HOITINK, HARRY A. J., 427
HOLCOMB, G. E., 28
HOLDEMAN, QUINTIN L., 373
HOLLAND, A. H., (531)
HOOKER, A. L., 708, (819), 1109
HORN, N. L., 293
HORTON, B. D., (620)
HOUGH, L. F., (682)
HRUSCHKA, HOWARD W., 219
HUGHES, JERRELL E., (474)
HUGHES, P. A., (328)
HUMPHREY, W., (940)
HUNTER, J. E., (694)
HUTTON, K. E., 776
HYRE, RUSSELL, 1131, (1134)

INMAN, R. E., 3
ISAKSON, O. W., (935)
IZADPANAH, K., (339)

JACKSON, L. W. R., 465
JACKSON, NOEL, 168
JACOBSEN, B. J., 456
JANSEN, WILLIAM P., 1053
JENSEN, J. J., (364)
JEYARAJAN, R., (617)
JHOOTY, J. S., 1049
JOHNSON, A. W., 952
JOHNSON, FOLKE, (268), (701)
JOHNSON, J. T., 979
JOHNSTON, H. WINSTON, 91
JONES, BILLY L., 603
JONES, ROBERT W., (822)
JOSHI, L. M., 391, 594

KABLE, P. F., (776)
KAHN, ROBERT P., 675
KAISER, WALTER J., 246
KAMAL, M., 241
KAPOORIA, R. G., 646
KARPEN, DANIEL N., 945
KAUFMAN, J., (263), (856)
KAVANAGH, JAMES A., 453
KEIL, H. L., 208
KENKNIGHT, GLENN E., 12
KESSLER, KENNETH J., Jr., 16
KHURANA, S. M. PAUL, 181, 437
KINGSOLVER, C. H., (130)
KINMAN, M. L., (111)
KITSOS, G. T., (561)
KLEIN, M., 735
KNESEK, J., 497
KNIGHT, ROBERT J., Jr., (569)
KNIGHTEN, J. L., (560)
KNORR, L. C., 630, 1092
KOIKE, H., (554)
KOONS, CAROL, (277)
KREBILL, R. G., 853
KUMMEROW, JOCHEN, (403)
KUNIMOTO, R. K., (694)
KUSHALAPPA, A. C., 788
KYROU, N. C., 134

LACY, M. L., (290)
LaFEVER, H. N., (156)
LAMBERTI, FRANCO, 717
LAMEY, H. A., 56, 931
LATHAM, A. J., 408, 725
LAUGHLIN, C. W., (287)
LEE, SOON HYUNG, (565)
LENZ, J. V., (1012)
LEONE, IDA A., (704)
LESEMANN, D., (851)
LEWIS, F. H., (217)
LEWIS, J. A., (114)
LIM, S. M., (708), (819), (1109)
LINDNER, R. C., (44)
LINDSEY, M. F., (281)
LINE, R. F., 992
LING, K. C., 565
LLOYD, EDWARD H., Jr., (387)
LOCKWOOD, J. L., 846, 849
LOEBENSTEIN, G., (643)
LONG, P. G., 93
LORING, L. B., 700
LOVREKOVICH, H., (713)
LOVREKOVICH, L., 713
LOWNSBERY, B. F., (920)
LUCAS, L. T., (976)
LUEDDERS, V. D., (1088)
LUHN, C. F., 1055
LUKENS, R. J., 152
LUKEZIC, F. L., (811)

MAAS, JOHN L., (362), 667, (739), 883, 887
MacSWAN, IAIN C., (887)
MAGGENTI, A. R., 1012, 1014, (1016)
MAHMOOD, TARIQ, 881
MAI, W. F., (368), 792
MAJORANA, G., 389
MANNING, W. J., 328
MANSOUR, KAMAL, (666)
MARCLEY, M. D., 1061
MARINOS, M. G., (706)
MARLATT, ROBERT B., 199, 569
MARSH, P. B., (408)
MARTIN, W. J., 1056
MARTINEZ, A. L., 1007
MASON, GARLAND N., (931)
MASSIE, L. B., (146)
MASTERNAK, ANNA, (81)

MASTERNAK, H., (640)
MAY, C., (858)
MAY, JAMES, (490), (493)
McCARTER, S. M., 586
McCRACKEN, F. I., 322
McDONNELL, P. F., 83
McGLOHON, NORMAN E., (12)
McLEAN, D. M., 372
McNEILL, K. E., 829
MELCHING, J. STANLEY, (130), 512
MENON, M. R., 34
MEYER, KATHRYN B., 899
MEYER, W. A., 506
MICKAIL, KAMEL Y., (666)
MILHOLLAND, R. D., (122), (125)
MILLER, C. C., (1041)
MILLER, H. N., 966
MILLER, J. R., (8)
MILLER, P. M., 25, 27, 261, 536, 672, 781, 995
MILLER, PAUL R., 1122, 1134
MILLER, V. L., (235), (377)
MILLIKAN, D. F., (81), (640)
MINK, G. I., (497)
MOELLER, C. F., (414)
MONCRIEF, J. B., (419)
MONROE, R. L., (675)
MOORE, K. J., (774)
MOORE, WILLIAM F., 1104
MORTON, H. L., 612
MUSSON, M. D., (1109)

NASH, G. T., (349), (927)
NAULT, L. R., 156
NEELY, DAN, (58), 127
NELSON, BARBARA V., (899)
NELSON, J., (187), (531), (689)
NELSON, R. R., (277), 1123
NEMEC, S., 49, 416
NETZER, D., 232, 909
NEWMAN, B. E., (28)
NICHOLLS, THOMAS H., 731
NIELSEN, L. W., (315), (979)
NOEGEL, K. A., (966)
NOME, SERGIO F., (939)
NORA, D. M., (1007)
NUQUE, F. L., (1045)

O'BANNON, J. H., 692
OITTO, W. A., (835)
ORELLANA, R. G., 111, 891
ORLOB, G. B., 1029
OSBORNE, W. E., (1064)
OSHIMA, NAGAYOSHI, (266)
OTTA, J. D., 332
OU, S. H., 1045

PALMER, J. G., (598), 858
PALMITER, D. H., 101, 105, 395
PAPAVIZAS, G. C., 114
PAPIA, P. M., (328)
PARACER, S. M., 920
PARISH, C. L., (44), (651)
PARKER, K. G., (792)
PARTRIDGE, A. D., 701
PATHAK, V. N., (976)
PATTON, R. F., (612)
PAULSEN, AVELINA Q., 627
PAULUS, A. O., 531, 689
PAYNE, J. A., (620)
PEACHER, PAUL H., (713), (931)
PEIRCE, L. C., 681
PÉREZ, J. ENRIQUE, 212
PERRY, ROBERT S., 451
PETERSEN, D. H., (1123)
PETERSON, GLENN W., 572
PETERSON, J. L., 606, 1037
PHELPS, WILLIAM R., 1031
PHILLIPS, D. V., 987
PINCKARD, J. A., (76), (524)
PITRE, H. N., (876)
POOL, R. A. F., 825
PORTER, D. MORRIS, 955
POWELL, DWIGHT, (832)
POWELSON, R. L., (992)
PRABHU, A. S., 959
PRASAD, KRISHNA, 40, 1026
PRATT, R. G., (583)
PRING, DARYL R., 550
PROVVIDENTI, R., 744
PUCHALA, Z., (640)
PURCIFULL, D. E., (161)
PYEATT, L., (940)

QUINIONES, S. S., 301

RAABE, R. D., (399)
RADEWALD, J. D., 187, 940
RATCLIFFE, T. J., (952)
RAWLINS, T. E., (741)
RAYCHAUDHURI, S. P., (34), (391), (594)
REINERT, R. A., 8
RENFRO, B. L., (391), (594), (989)
REYNOLDS, JAMES E., 223
RHOADES, H. L., 411, 635
RHODES, A. M., (896)
RICH, A. E., (360)
RICKARD, S. F., 414, 1041
RIEDEL, R. M., 251
RIFFLE, JERRY W., 752
RIGGS, R. D., (544)
RIVES, MAX, 915
RIZK, ZAKI, 666
ROHRBACH, K. G., 694
ROMIG, R. W., (256)
RONCADORI, R. W., (586)
ROTEM, J., 768
ROTHMAN, P. G., (256)
ROWE, C., (501)
RUPPEL, E. G., 176
RUSH, M. C., 342

SAAD, A. T., (435)
SAARI, E. E., (391), (594), (635)
SAENGER, H. L., 136
SAFFARIAN, A., (339)
SANCHEZ A., LUIS, (542)
SANDERS, H., (49)
SANDHU, R. S., (337)
SAYDAM, C., (241)
SCHARPF, R. F., 275
SCHEIFELE, G. L., (277), 501
SCHENCK, N. C., 1127
SCHLÖSSER, W. E., (435)
SCHMITTHENNER, A. F., (427)
SCHREIBER, LAWRENCE R., 240
SCHROEDER, W. T., (744)
SCHUSTER, JAMES E., 58
SCHUSTER, M. L., (557)
SCHWARZ, R. E., 1003
SCOTT, D. H., 362, (739)
SCOTT, GENE E., (307), 483
SCOTT, O. J., (498)
SELISKAR, CARL E., (844)
SEMENIUK, P., 598
SEMLER, LARRY F., (63)
SEQUEIRA, LUIS, 754
SHAFFER, W. H., Jr., 203
SHAH, S. MOIN, (1092)
SHANE, JAMES F., (723)
SHARAFEH, M., (339)
SHARP, E. L., (992)
SHARVELLE, ERIC G., 1112
SHAUNAK, K. K., 876
SHERWOOD, CHARLES H., (1026)
SHIBUYA, F., (187), (531), (689), (940)
SHORTLE, W. C., 360
SHRUM, ROBERT D., (512)
SIDDIQUI, I. A., 972
SILL, WEBSTER H., Jr., (627)
SILVERBORG, SAVEL B., (462)
SIMONS, M. D., 807
SIMPSON, M. E., (408)
SINCLAIR, J. B., (506)
SINGH, RAGHBIR, (617)
SINHA, M. K., 617
SKELLY, J. M., (811)
SKILLING, DARROLL D., 132, 663, (731)
SMITH, A. M., 774
SMITH, C. B., (141)
SMITH, D. H., 217
SMITH, D. R., (708), 819, (1109)
SMITH, N. A., (849)
SMITH, P. G., 786
SMITH, RICHARD S., Jr., 872
SMITH, WILLIAM H., 1021
SMITHSON, HARRIET, (700)
SOWELL, GROVER, Jr., (60), (880)
SPALDING, D. H., 655
SREERAMULU, T., 226
SRIDHAR, R., 272, 632
STACE-SMITH, R., (701)
STALEY, JOHN, (275)
STAMBAUGH, W. J., (560), (799)
STAPLES, ROBERT, (1053)
STATLER, GLEN, D., 698
STEWART, D. M., 256
STIENSTRA, WARD C., 290
STOKES, D. E., 287
STONER, W. N., (191)
STRIDER, D. L., 36, 976
STYER, W. E., (156)
STYNES, B. A., (774)
SUNDARAM, S., (1035)
SUTHERLAND, JACK R., 165

TAKACS, D. J., 620
TAKAHASHI, Y., (1029)
TAYLOR, D. P., 435, (972)
TAYLOR, G. S., (672)
TAYLOR, JACK, 871, 1039
TEDDERS, W. L., (620)
TEWARI, J. P., (727)
THANASSOULOPOULOS, C. C., 561
THAUNG, MAUNG MYA, 427
THOMAS, C. A., 300, 1010
THOMAS, CLAUDE E., 108, 155
THORNBERRY, H. H., 67
TIPPETT, R. L., 554
TOLIN, SUE A., 935
TOMLINSON, J. A., 150, (370), (734)
TORRES, H., 315
TRAYLOR, JACK A., (822)
TROUTMAN, J. L., (935)
TU, CHIN-CHYU, 29
TU, J. C., 384

ULLSTRUP, A. J., 630, 658, 1100, 1103

Van ARSDEL, E. P., 669
van der ZWET, T., 96, 835
VARGAS, J. M., Jr., 1075
VARNEY, E. H., (682)
VITTAL, B. P. R., (226)
VOLCANI, ZAFRIRA, 804
VÖRÖS, Jr., (286)

WADDELL, CLINTON D., (663)
WAGNON, H. KEITH, (822), (825)
WALKER, D. D., 1114
WALKEY, D. G. A., 370, 802
WALLACE, HUEY N., 713
WALLIN, JACK R., (959), 1129, (1134)
WALTON, G. S., (536)
WARD, H. N., (701)
WATERS, E. C., (783)
WATERWORTH, H. E., 1059
WATSON, D. R. W., 86, 1068
WEATHERS, L. G., 542
WEAVER, L. O., (208)
WEBB, R. E., (155)
WEBER, RAY, (1031)
WEHLBURG, C., 827
WEHUNT, E. J., (620)

WEIGLE, JACK L., (40), (1026)
WELLMAN, FREDERICK L., 355, 539
WELLS, JOHN M., 431
WELSH, MAURICE F., 490, 493
WHEELER, J. E., (179)
WHIDDEN, R., (907)
WHITEHEAD, W., (501)
WHITESIDE, J. O., 608, 865
WHITTINGHAM-JONES, SYLVIA G., (802)
WIESE, M. V., 635
WILCOXSON, R. D., (391), (594)
WILES, A. B., (508)
WILLIAMS, HAROLD E., 822, (825)
WILLIAMS, L. E., (156)
WILLIAMS, PAUL H., (453), (456)
WILLIAMS, R. E., (996)
WILSON, CHARLES L., 905
WILSON, G. F., 896
WINSTEAD, E. E., (243)
WOLFE, R. D., 471
WONG, T. K., 368
WOOD, R. E., (330), 438
WORF, G. L., (281)
WYLLIE, T. D., (1088)

YARWOOD, C. E., (399)
YASSIN, A. M., 528
YODER, D. L., (846), (849)
YOUNGSON, C. R., 196

ZEHR, ELDON I., 516
ZETTLER, F. W., (161), (893)
ZIMMER, D. E., 364
ZUCKER, MILTON, 863
ZUMMO, NATALE, 723
ZUTRA, D., (804)

SUBJECT INDEX

Abaca (Musa textilis): Pseudomonas solanacearum isolated from abaca in the Philippines but of low virulence, 516

Abies bracteata (bristlecone fir): first known disease of North American tree species, caused by needle cast fungus, Lirula nervisequia var. conspicua, in California, 275

--- concolor (white fir): host of true mistletoes (Phoradendron spp.); both hyper- and autoparasitism observed in Phoradendron species, 15; Pleospora sp. associated with cankers on curled twigs at three locations in northwestern Oregon, 700; control of Fomes annosus with borax, 872

--- grandis (giant fir): twig dieback, caused by Pleospora sp., in Oregon, 700

Aceria tulipae: 156; 635

Additives: for tree cavity fills, burlap, and fungicides added to urethane foam, 858

Adjuvant: six surfactants tested as adjuvants for fungicidal action of benomyl against Verticillium on inoculated cotton plants, 741

Ageratum: stem rot, caused by Alternaria zinniae, found for first time in the United States in Illinois commercial greenhouse, 1095

Air pollution: field studies of air pollution injury to vegetation in Cincinnati, Ohio, 8; toxicity of sulfur dioxide to highbush blueberry, 704; effect of ozone and sulfur on ponderosa pine geographic race plantation in Washington, 945; host response among different Phaseolus vulgaris cultivars and accessions shows variation, 1026

Alabama: 66; 322; 408; 725

Alfalfa: seedlings tested for sensitivity to high levels of air pollution, 8; production of plant parasitic nematodes on callused host tissue, 251; virus-infected seeds in alfalfa seed lots, 591; pathogenicity of Sclerotinia trifoliorum to alfalfa and crownvetch, 811

Almond (see also Prunus): seed transmission of necrotic ring spot virus, 822

Alternaria alternata: 453

--- circinans: 28

--- tenuis: 586; 647; 655; 796; 871

--- triticina: 594

Anguina tritici: 594

Announcements: for persons on free mailing list, 272; confirmation of occurrence of coffee rust, caused by Hemileia vastatrix, in Brazil, 355; for rapid interchange of bacterial cultures, 814; validation of plant introduction (PI) numbers in manuscripts for Plant Disease Reporter, 912

Anthracnose: of tomato, 121 accessions evaluated for resistance, PI 272636 showed only trace amounts of infection in the field, 32; of Sorghum bicolor, caused by Colletotrichum graminicola; over 300 of 779 introductions screened exhibited good field resistance, 60; of poppy, caused by Gloeosporium cingulata, not previously reported on oriental poppy, 1037

Antibiotics (see also Chemicals and Fungicides): cycloheximide: applied directly in fuel oil to Hypoxylon cankers on aspen arrested canker activity for 5 years and tree survival increased with increase in concentration of antibiotic, 462; as a dip effective against powdery mildew (Sphaerotheca pannosa)

on rose seedlings, 598; effective against Ceratocystis fimbriata infection of Hevea brasiliensis, 897; cycloheximide acetate and cycloheximide thiosemicarbazone as spray are distributed systemically in 1- and 6-year old Jonathan apple trees, 996; gave most consistent effective results for suppression of white pine blister rust cankers in tests with different carriers, 1031
---: oxytetracycline, control of peach bacterial spot requires frequent applications of antibiotic, 208
---: streptomycin, in combination with fungicide and insecticide, gave better control of fire blight of apple than streptomycin alone, 203
---: tetracycline hydrochloride, (achromycin and tetrachel) suppressed Philippine greening disease of citrus, 1007
Aphanomyces euteiches: 943
Aphis craccivora: 344; 935
--- gossypii: 437
Apocarpy: of papaya fruits, caused by distortion ringspot virus, 181
Apple: Botryosphaeria ribis, isolates pathogenic to 'Golden Delicious' apples after eight years' storage under sterile mineral oil at 2-5°C, 217; fruit blotch disorders (7) on six apple cultivars in British Columbia, two on 'Delicious' apple were graft-transmitted, 490; preplant nematicide treatment with DD-MENCS resulted in greater apple tree growth than in untreated plots, and in better root systems, 792; three systemic fungicides tested for distribution in sprayed 'Jonathan' apple trees for concentration and duration of chemical in fruit, stem and roots, 996
---: fire blight: caused by Erwinia amylovora, controlled best with night sprays consisting of steptromycin, sulfur and glyodin, 203; twig blight, presence and movement in inoculated 1-year-old apple trees, 576; severity on 'Mutsu' apple trees, 638; severe on Mollie's Delicious cultivar in Georgia, 1039
---: little leaf, a probable temperature-induced genetic disorder, 1059
---: postharvest decay, of fruit by Nectria galligena curbed with benomyl spray or dip, 83
---: powdery mildew, Podosphaera leucotricha controlled with benomyl, 81
---: scab: spring application of benomyl to overwintering apple leaves reduced discharge of ascospores 93%, thiabendazole and urea less effective, 27; distribution of scab, caused by Venturia inaequalis, in the State of Washington, 63
---: virus: pustule canker and freckle scurf, two graft-transmissible bark diseases of apple are described, 44; host range comparisons indicate a possible relationship between green mottle, chat fruit, and flat apple viruses, 105; fruit wrinkle, a graft-transmissible abnormality of 'Newtown' apple in British Columbia, 493
Arachis hypogaea (see Peanut)
Arceuthobium campylopodum f. blumeri: 488
--- --- f. cyanocarpum: 441
--- --- f. tsugensis: 488
Argentina: 722; 939
Arizona: 176; 179; 1064
Arkansas: 471
Arundinaria gigantea (bamboo): sugarcane mosaic virus identified, possibly first record of a virus disease of bamboo, 815
Ascochyta gossypii: 324
--- rabiei: 442
Aspen (see Populus)
Aspergillus spp.: 796
--- flavus: 408
--- niger: 337
Aspirator: compressed air aspirator for use in collection of large numbers of viruliferous plant hoppers, 56
Assay: of fungicides against Helminthosporium papulosum f. pyri, cause of blister canker of pear; maneb, ferbam and thiram showed greatest inhibition of fungus, 296
Australia: 622; 774; 776; 903; 1061
Autoecious rust: western gall rust, Peridermium harknessii, 853
Autoparasitism: in true mistletoes, as well as hyperparasitism, found in Phoradendron species in California, 15

Bacillus subtilis: 446
Bacterial diseases: of abaca and banana, isolation of Pseudomonas solanacearum in the Philippines, 516; Pseudomonas syringae caused leaf and stem lesions on Austrian winter pea, a new host for the bacterium, 467
---: bacterial blight: of dry bean, caused by Xanthomonas phaseoli, Jules variety very tolerant; registered and certified seed of this variety will be produced in 1971 and 1972 in Nebraska, 557; fuscous and bean common blight first found in New Zealand in 1970, caused by imported seed, 1068; of cotton, by Xanthomonas malvacearum, 324; ammonia is apparently involved in formation of cotton blight symptoms and in development of the hypersensitive necrotic reaction caused by Pseudo-

monas tabaci in cotton, 713; of guar, new virulent race found in Texas, 111; of soybean, reported from New Zealand for the first time, 86; of statice, in Florida, 759
---: bacterial canker: of French prune, Pseudomonas syringae isolated from cankers every month of the year in California, pruning wounds are potential sites of entry, 332; of tomato, root and stem inoculations of tomato seedlings compared as well as age of inoculum, 36; variation in virulence of Corynebacterium michiganense, 796
---: bacterial scab, of gladiolus, caused by Pseudomonas marginata, was less severe on benomyl-treated corms than with thiram treatment, 289
---: bacterial soft rot: of lettuce, a serious market disease caused by Erwinia carotovora and Pseudomonas sp., 263; caused by P. marginalis, P. cichorii, and E. carotovora, 964
---: bacterial spot: of peach, oxytetracycline as weekly spray gave control of Xanthomonas pruni in orchards in Maryland, 208; cultivars, Redhaven, Oriole, Raritan Rose, Royalvee, Loring, and Veteran found to have some resistance to bacterial spot and to Valsa canker, in New York tests, 395
---: bark canker, of Persian walnut, in the central valley of California, 373
---: black leg (Erwinia atroseptica) and brown rot (Pseudomonas solanacearum) of potato reported for the first time from Peru, 315
---: fire blight: of pear, caused by Erwinia amylovora, seedling screening procedure, 96; controlled best on apple with combination spray of streptomycin, sulfur and glyodin, in Missouri, 203; stem apex inoculation of girdled apple trees showed E. amylovora moves downward in stems exclusively in the phloem, 576; on young 'Mutsu' apple trees, 638; system for estimating severity of fire blight in pear, 835; of Mollie's Delicious apple cultivar, 1039
---: phloem canker, of Persian walnut, caused by Erwinia rubrifaciens, is serious disease in central valley of California because one-fourth of walnut acreage is planted to the highly susceptible Hartley variety, 373
Bamboo (see Arundinaria gigantea)
Banana: stem-end rot disease, caused by Gloeosporium musarum, 93; vascular wilt of banana, caused by Pseudomonas solanacearum, found in Philippines, 516
Barley: barley stripe mosaic virus, field spread in Montana greater in 1968 than 1967 but there was no corresponding increase in percentage infection of diseased plots, 749; brown loose smut (Ustilago nuda) of barley, believed to be first report on leaves of Larker cultivar, 927
Basidiomycete: unidentified Basidiomycete appeared on bentgrass turf after use of benomyl in Australia, 774
Bean: Aphanomyces euteiches races identified by use of differential bean series, 943; bell bean (Vicia faba var. minor), most reliable assay host for fizzletop virus of pea, 355; broad bean, Helminthosporium spiciferum, cause of severe root and stem discoloration of broad bean, the first record of this fungus in the Sudan and possibly first record of pathogenicity to broad bean in other parts of the world, 405; mung bean (Phaseolus aureus), new record of leaf spot, caused by Alternaria alternata, in India, 453; moth bean (Phaseolus aconitifolius), Colletotrichum truncatum, cause of new leaf spot in India, 976
---: air pollution: injury to bean in Cincinnati, Ohio, 8; (Phaseolus vulgaris), variation in ozone phytotoxicity among bean cultivars, 1026
---: blight: Great Northern dry bean variety Jules is highly tolerant of Xanthomonas phaseoli, cause of common blight, and shows promise for use in commercial bean seed production in Nebraska, 557; common blight and fuscous blight of dwarf bean in New Zealand in 1970, 1068
---: nematodes: pole, foliar spray of DuPont's 1410, nematicide - insecticide controlled Pratylenchus scribneri, 187; newly found Rotylenchulus reniformis in Lebanon could be threat to vegetable production in coastal region, 435
---: Rhizoctonia: damping-off resistance, additional sources of resistance and screening techniques based on host reaction are reported, 40; Rhizoctonia stem canker on mung bean (Phaseolus aureus) in Iran, 246
---: root rot: caused by Thielaviopsis basicola, several soil amendments reduced root rot of bean in the greenhouse, 114; root rot caused by Fusarium solani f. phaseoli, bean varieties with purple hypocotyls and black seeds were more resistant to fungus than those lacking this color, 698
---: summer death disease, field reactions

of 61 bean varieties and breeding lines in Australia, 903
---: virus, peanut stunt virus caused disease of epiphytotic proportion to beans in North Carolina, 183
Belonolaimus longicaudatus: 635
Bemisia tabaci: 34; 528
Bioassay: studies on distribution and length of detection of benomyl in stem sections and leaf tips of tomato inoculated with Verticillium albo-atrum and Penicillium expansum, 12; of cantaloup leaf discs, correlation shown between powdery mildew incidence and the amount of benomyl in plants after root application of fungicide, 232; persistence and distribution of 8-quinolinol phosphate, cycloheximide acetate, and cycloheximide thiosemicarbazone in 1- and 6-year-old apple trees by Cytospora bioassay, in Idaho, 996
Biological control: of Fusarium stem-rot of carnation cuttings, by infesting rooting media with Bacillus subtilis, 446
Birch (Betula alleghaniensis): Diaporthe alleghaniensis and Valsa ambiens found associated with canker and dieback of yellow birch seedlings, 16
Black leg: of potato, caused by Erwinia atroseptica in Peru, 315
Blast disease: of rice: 11 races of Pyricularia oryzae identified in Nigeria, 520; excessive application of nitrogenous fertilizers found to increase susceptibility of rice to blast, 632; on rice seed in the United States, 931; classification of many different races from single conidial isolates, 1045
Blight: Ascochyta, of cotton, estimate of loss by State in 1969, 324
---: bacterial: of dry beans, Jules variety found to be highly tolerant to Xanthomonas phaseoli in Nebraska tests, 557; caused by X. phaseoli and X. phaseoli var. fuscans, from imported seeds, 1068; of cotton, estimate of cotton loss in 1969, 324; increase of ammonia in detached cotton leaves after inoculation with Pseudomonas tabaci or Xanthomonas malvacearum, 713; of guar, caused by Xanthomonas cyamopsidis, 111; of soybean, caused by Pseudomonas glycinea recorded for first time from New Zealand, 86
---: Botrytis, of hydrangea, benomyl gave excellent control in greenhouse tests, 606
---: fire: of apple, four night sprays of combination antibiotic, fungicide and insecticide better than antiobiotic alone in control of twig blight caused by Erwinia amylovora, 203; movement and persistence of E. amylovora in shoot, root and stem of apple, 576; Mutsu apple variety growing on EM26 and MM 106 rootstocks more severely infected than Delicious and McIntosh varieties in Vermont test, 638; Mollie's Delicious cultivar very susceptible in Georgia, 1039; of pear, needle injection and stem decapitation better than spray inoculation for screening of pear seedlings for resistance, 96; number of twig infections, age of wood infected, and percent of pear tree blighted are basis for scoring system for severity of infection, 835
---: Fusarium blight: of 'Merion' Kentucky bluegrass, chemical control best with benomyl spray in spring, 771; least infection from F. roseum observed under conditions of low fertility and high moisture availability, 1018.
---: late blight: of potato, where macroclimate favors plant disease development, the microclimate of the field is of minor importance, 768; of tomato, caused by Phytophthora infestans, detached leaf method for screening tomato plants for single gene resistance to race O, 681
---: leaf, of wheat, caused by Alternaria triticina, 594
---: ---: northern corn leaf blight, chlorotic-lesion type of resistance, 896
---: ---: southern corn leaf blight: inbred lines and hybrids with Texas male-sterile cytoplasm show increased susceptibility to Helminthosporium maydis, 501; reaction of inbred lines of corn with T or P cytoplasm, 708; physiologic races of pathogen, 819; special issue on occurrence of epiphytotic in the U.S., from H. maydis, 1099-1136
---: ---: yellow leaf blight: caused by Phyllosticta sp., data suggest that male-sterile cytoplasm usually increases susceptibility of maize inbreds and single crosses to the disease, 277; of corn in Wisconsin, its history and the reactions of inbreds and crosses to the fungus, 281
---: Macrophoma and Septoria blights of potato found in Peru, 315
---: needle: red band needle blight, infection of Monterey pine graftings with Dothistroma pini in Chile, 403; of Austrian pine, benomyl + polyvinyl chloride, Bordeaux and TC-90 effective in control, 679
---: seedling blight: of corn, caused by Fusarium moniliforme, 307; response of corn seedlings to F. moniliforme

and a toxic material extracted from the fungus, 483
---: Typhula blight of bentgrass in Michigan controlled with chloroneb, 1075
Blueberry: eradication of Botryosphaeria corticis from propagation of wood of Vaccinium corymbosum with fungicidal dip of BSM-11, 122; V. australe, four cultivars tested showed variability in response to sulfur dioxide as an air pollutant, and were rated as fairly tolerant, 704
Blue mold: of stored pears, controlled with postharvest treatment of benomyl or thiabendazole, 655
Boerhaavia diffusa: weed host of brinjal mosaic disease in India; virus found to be a new strain of cucumber mosaic virus 1, 437
Bolivia: 675
Boll rot (see Rot)
Botryodiplodia theobromae: 809
Botryosphaeria corticis: 122
--- ribis: 217
Botrytis cinerea: 132; 263; 431; 606; 647; 694; 883; 964
Brazil: 355; 539
Brinjal (see Eggplant)
Brinjal mosaic disease: 437
Bristlecone fir (see Abies bracteata)
Britain (see England)
British Columbia (see Canada)
Brown patch: of turfgrass, caused by Rhizoctonia solani, fungicidal control, 1080
Brown rot: caused by Sclerotinia laxa, found on grape, in New Zealand, 504
---: of potato, caused by Pseudomonas solanacearum in Peru, 315
Brown stain: of head lettuce, a disorder found in western head lettuce, 856
Bulb mite (Rhizoglyphus echinopus): control of bulb mite that transmits bacterial scab of gladiolus, benomyl more effective than thiram, 289
Bunch disease: of pecan, in Georgia, first report of disease east of Meridian, Mississippi until this discovery, 12
Bunt: common bunt, caused by Tilletia caries, and T. foetida, in India, 391
---: karnal bunt, caused by Neovossia indica, in India, 391
Burma: 427

Cactus (see also Zygocactus truncatus): control of cactus cyst nematodes on Christmas cactus was best with BAY 25141 and V-C 9-104 692; witches'-broom disease of Opuntia tuna, cactus virus X found in cactus grown as house plant, 851
Caladium bicolor: root-knot nematodes, nematicide treatments compared to hot water treatments for control of Meloidogyne spp. in tubers, 411
California: 15; 91; 275; 332; 373; 399; 488; 531; 786; 872; 920; 940; 964; 969; 1014; 1055
Canada: 91; 165; 490; 493
Canker: of yellow birch seedlings, Diaporthe alleghaniensis and Valsa ambiens were isolated from some diseased stems, 16
---: bacterial: of tomato, seedling inoculations with Corynebacterium michiganense, 36; of French prune, caused by Pseudomonas syringae, 332; bark canker and phloem canker of Persian walnut in California, 373
---: blister, of pear, bark disease of pear trees found in Massachusetts in 1966, caused by Helminthosporium papulosum f. pyri, 296
---: of cocoa, caused by Phytophthora palmivora by stem inoculation and natural infection, cocoa cultivars were assessed for susceptibility, 1035
---: Cytospora: of pecan, found in Arizona, first known published report of disease on pecans in the United States, 179; of peach in Illinois, 832
---: Endothia parasitica pathogenic to chinkapin in Louisiana, 713
---: Hypoxylon cankers on aspen arrested after direct applications of cycloheximide, in New York, 462
---: Pezicula, of apple, controlled with benomyl and Cercobin in Poland, 640
---: pustule canker suggested as name for graft-transmissible bark disease of 'Red Delicious' apple, 44
---: Rhizoctonia stem canker of mungbean in Iran, 246
---: Scleroderris canker: found on red and jack pine in Minnesota in July, 1969, 132; on red pine, controlled with maneb or ziram on nursery stock, 663
---: stem canker of blueberry, caused by Botryosphaeria corticis, eradicated from dormant cutting wood with fungicide BSM-11, 122
---: Valsa canker of peach, relative resistance of 26 cultivars to bacterial spot and Valsa canker, 395
Carnation: cuttings treated with Bacillus subtilis incorporated into rooting medium had reduced Fusarium stem-rot infection, also developed more extensive root systems, 446
Carotenoid content: pigment content of leaves of Chenopodium nuttalia reduced when infected by Peronospora farinosa, 762
Carriers: chemical carriers Mobil Sol L,

No. 2 fuel oil, vegetable oil and xylene effective for chemotherapeutants on scarified cankers for control of blister rust on eastern white pine, 1031
Cassava mosaic virus: 34
Cavariella aegopodii: 370
--- pastinacae: 370
Cedar posts: growth of shade tobacco was greater around newly set cedar posts than pine posts in Connecticut, 536
Celery: western celery mosaic virus isolated from celery, coriander and parsley, in Britain, 370; Septoria leafspot controlled with several fungicides on a 14-day preventive spray schedule, 531; seed transmission of strawberry latent ringspot virus in celery, 802
Cenangium singulare: 460
Cephalosporium spp. (= Phialophora obscura): 669
--- acremonium: 989
--- gregatum: 987
Ceratocystis alba: 460
--- crassivaginata: 460
--- fimbriata: 897
--- populina: 460
--- tremulo-aurea: 460
--- ulmi: 58; 127; 240; 266; found in Wyoming for first time in summer of 1969, 547; 701; 706; found in Texas for the first time, 931; 985
Cerotoma ruficornis: 212
--- trifurcata: 1053
Ceratostomella (Ceratocystis) merolinensis: 905
Cercospora sp.: 199
--- arachidicola: 955
--- insulana: 759; 760
--- nicotianae: 666
--- oryzae: 272
--- personata: 955
--- salicina: 939
Chalaropsis thielavioides: 451
Charcoal rot (see Rot)
Chat fruit (see Apple virus)
Chemicals: role of ammonia in formation of cotton blight symptoms in cotton leaves from bacterial blight infection by Pseudomonas tabaci, 713
---: borax, dry powder most effective in control of Fomes annosus infection of white fir stumps, in California, 872
---: EDTA (ethylenediaminetetraacetic acid) application to potato plugs decreased soft rot from Erwinia carotovora or Pseudomonas fluorescens, 863
---: herbicides: sodium cacodylate and picloram not effective in prevention of Fomes annosus infection when used to thin young stands of western hemlock in Washington, 330; potassium azide shows fungicidal qualities as a seed treatmen 604
---: indole butyric acid (IBA) a rooting hormone used with benomyl or thiabendazole on rhododendron cuttings, 427
---: insecticides: Dasanit (BAY-25141) and ethion with thiram plus a thiram seed treatment controlled onion smut and onion maggot in field-seeded onion at three planting depths in Michigan, 290; EPN (O-ethyl O-p-nitrophenyl phenylphosphonothioate) combined with fungicides effective in reduction of cotton boll black rot in Taiwan, 29; glyodin, with streptomycin and sulfur, as a spray for control of fire blight of apple, 203; parathion in greenhouse fumigation found to affect the expressi of crown rust on oat cultivars, 807
---: nitrogenous fertilizers, at high rates increased susceptibility of rice to blas disease, 632
---: 8-quinolinol phosphate, cycloheximid acetate, and cycloheximide thiosemicarbazone, systemic distribution in sprayed 'Jonathan' apple trees in Idaho, 996
---: Rindite (material used for breaking th rest period of potatoes) found to be effective in control of encysted golden nematode larvae, 306
---: salt: relative sodium chloride tolerance of common roadside trees in southeastern New Hampshire, 360; salt (calcium and sodium) damage to eastern white pine in Connecticut from winter-salting of an interstate highway 1021
---: sodium salt of dehydroacetic acid dip for control of postharvest decay of strawberries compared with modified atmosphere and heat treatments, 431
---: sulfur dioxide as an air pollutant, highbush blueberry cultivars were fair tolerant to exposure in experimental chambers, 704
---: sulfuric acid, for damping-off control of pine seedlings was no advantage if bed had been previously fumigated with methyl bromide, 572
Chenopodium nuttalia: changes in chlorophyll and carotenoid content in C. nuttalia leaves by Peronospora farinosa, 762
Cherry (see also Prunus): occurrence of short stem virus of sweet cherry reported from Oregon, 651; top- or branch-killing of inoculated Kwanzan flowering cherry nursery trees caused only by some strains of the Prunus

ringspot virus, not by 11 other Prunus viruses tested, 1073
Chickpea (Cicer arietinum): a limiting factor in high chickpea yields in Israel is infection by Ascochyta rabiei, 442
Chickweed: seed transmission of cucumber mosaic virus from infected chickweed plants averages 5 to 8%, 150
Chile: 403; tristeza disease of citrus found in trees imported to Chile from the U.S. over 20 years ago, 542
Chinese gooseberry (Actinidia chinensis): stem-end rot, Phomopsis sp. identified as incitant on ripe fruit on Chicago market, 422
Chinkapin (Castanea pumila): cankers of Endothia parasitica infection found for the first time on chinkapin in Louisiana, 713
Chlorophyll: Graphiola leaf spot infection of date palm resulted in reduced content of chlorophylls a and b and of total chlorophyll, 617; reduction of chlorophyll and carotenoid pigments in Chenopodium nuttalia leaves by Peronospora farinosa, 762
Chlorotic-lesion resistance: to Helminthosporium turcicum, 896
Chrysanthemum: Rhizoctonia stem rot, thiram-captan drench, developed for Pythium wilt, effective against R. solani at all stages of disease, 679
Citrus: a multiple sprouting disease in Southern Africa, 1003
---: citrus brown rot, factors contributing to the restricted occurrence in Florida, the predominant fungus involved was Phytophthora citrophthora, 608
---: citrus greasy spot, effect of foliar applications of fungicides on perithecial development of Mycosphaerella sp. in Florida, 865
---: Gloeosporium rot of 'Temple' oranges in Florida developed where fungicidal sprays were omitted, a new disease to Florida, 173
---: Geotrichum candidum, cause of sour rot of lemon in Turkey, 881
---: nematodes: decline, associated with presence of plant-parasitic nematodes in Iran, 339; Paratylenchus hamatus found associated with 60 declined citrus trees in Iran, apparently first report of this nematode in Iran, 1085
---: postharvest decay, orange, green mold control in artificially inoculated fruit, in relation to application time, 325
---: twig dieback of tangerine, benomyl sprays reduced incidence in Florida tests, 869
---: virus: susceptibility of sour orange, West Indian lime, common mandarin, lemon and sweet orange to tobacco necrosis virus, 389; tristeza virus found in Chile on trees imported in 1942 and 1947 from the United States, 542; greening virus disease in West Pakistan, 630; impietratura virus in Israel, leaf-flecking on indicator seedlings, associated with citrus impietratura, a possible indexing method, 643; double-bladed knife for making leaf-tissue grafts to transmit viruses, 907; citrus greening disease, in Philippines, suppressed with tetracycline antibiotics, 1007; greening disease of citrus in Nepal, 1092
Cladosporium herbarum: 647
Clasterosporium sp: 849
Claviceps purpurea: 844
Clitocybe tabescens: 620
Clover (Trifolium incarnatum): damping-off of crimson clover reduced by 5-minute treatment with radio-frequency energy (rf), 268; peanut stunt virus, an overwintering host in white clover, in Virginia, 935
Clubroot: of cabbage seedlings, common under greenhouse conditions, 453; of cabbage, controlled with benomyl in greenhouse tests, 456
Coccomyces hiemalis: 81
Cochliobolus heterostrophus (Helminthosporium maydis): perithecia on corn leaves in Florida in fall, 1970, 1127
Cocoa: cocoa canker, caused by Phytophthora palmivora, cultivars assessed in Fiji for susceptibility, 1035
Coffee: announcement of coffee rust in Brazil, 355; coffee rust, caused by Hemileia vastatrix, is positively established in Brazil, 539
Colletotrichum circinans: first record in Argentina on onions, 939
--- coccodes: 32
--- falcatum: 226; 427
--- graminicola: 60
--- truncatum: 976
Colombia: 675
Colorado: 266
Commercial disease control: of stem-end rot of banana, benomyl, tested under commercial conditions, gave better control than thiabendazole, in Samoa, 93
Computer program: program available for use with electric psychrometer that is especially suited for sensing temperature and humidity at different heights, at widely spaced locations and in confined places, 19
Connecticut: 171; 261; 536; 672; 995; 1021

Coriander (Coriandrum sativum): host of western celery mosaic virus, 370
Corky root: of Douglas-fir (Pseudotsuga menziesii), relationship of Xiphinema bakeri to disease, 165
Corky root rot (see Rot)
Corn: methods for inoculating corn ears with Gibberella zeae and Diplodia maydis, 658
---: Cephalosporium stalk rot, root injury increased with increase in plant population and flooding, 989
---: Diplodia stalk rot, 900 plant introductions rated for resistance and 42 showed some degree of resistance, 624
---: northern corn leaf blight, chlorotic-lesion resistance to Helminthosporium turcicum in a sweet corn background, 896
---: Phyllosticta sp.: inbred lines show resistance to yellow leaf blight, but evidence indicates resistance may be lost upon conversion to Texas male-sterile cytoplasm, 277; inbreds and hybrids with Texas male-sterile cytoplasm showed increased susceptibility to yellow leaf blight and differences in reaction of inbreds and crosses indicate genetic control, 281
---: rust. caused by Puccinia sorghi, corn varieties tested in two locations in India for susceptibility, 'Cuzco' showed no infection, 788
---: seedling disease: caused by Fusarium moniliforme, 69 inbred lines and 14 single crosses evaluated for their reaction to inoculation with the pathogen; 10 inbreds classified as resistant, the rest as intermediate or susceptible, 307; seedlings inoculated with toxin produced by Fusarium moniliforme reacted similarly to those inoculated with fungus, with retardation in root growth, 483
---: southern leaf blight: cytoplasm inheritance of susceptibility to Helminthosporium maydis indicated in susceptible inbred lines and hybrids with Texas male-sterile cytoplasm, 501; susceptibility of inbred lines carrying the T or P cytoplasm for male sterility to H. maydis, 708; two physiologic races, Race T and O differ in pathogenicity on inbred corn seedlings, 819; a special issue devoted to this disease, to the physiologic races of the pathogen, and efforts made to avert or minimize a recurrence of the epiphytotic of 1970, 1099-1136
Coronilla varia (see Crownvetch)
Correction: (Vol. 53, No. 10, Page 785), 352
---: (Vol. 53, No. 11, Page 859), 448
---: (Vol. 53, No. 12, Page 947), 88
---: (Vol. 53, No. 12, Page 968), 536
---: (Vol. 54, No. 2, Page 91), 271
---: (Vol. 54, No. 3, Page 193), 535
---: (Vol. 54, No. 7, Page 545), 816
Corticium fuciforme: 1080
Corylus avellana (Filbert): a decline of filbert trees in Oregon of unknown origin, "filbert stunt" is proposed name for disorder, 69
Corynebacterium michiganense: 36; found for first time to be causal agent of leaf and spot disease of sweet pepper under field conditions in Israel, 804; 976
Corynespora cassiicola: 52
Cotton: estimated reduction in cotton yield as result of parasitic diseases, 324; ammonia as a symptom causing factor in bacterial blight of cotton, 713; six surfactants tested for adjuvant properties with benomyl against Verticillium albo-atrum, 741
---: boll rot: cotton boll black rot, caused by Diplodia gossypina, control with fungicide plus an insecticide, 29; Fusarium spp., Verticillium nigrescens, and Alternaria tenuis were fungi most commonly isolated from decaying bolls; primary means of entry was through the bracts, 586; eight cotton varieties evaluated for internal infection through various modes of entry to boll; nectariless character seemed to contribute to less total visible internal infection, 719; fungi associated with boll rot in the Yazoo-Mississippi Delta, 796
---: damping-off: caused by Rhizoctonia solani, effectively controlled with benomyl, 76; control of R. solani on cotton seedlings with a 1,4-oxathiin derivative, 524
---: fluorescence, internal fluorescent material found in stems and roots of cotton from Arizona is distinct from fluorescing BGY material associated with Aspergillus flavus infection in the fiber, 408
---: Verticillium wilt, differences in host reaction to fungus by cultivars and weeds indicates importance of non-host crops in a rotation to reduce inoculum potential, 508
---: virus: tobacco ringspot virus caused local and systemic symptoms on Coker 201 cotton variety when inoculated into cotyledons, 342; transmission of leaf curl virus by single insects of Bemisia tabaci, 528
Cowpea (Vigna sinensis)(Southern pea): powdery mildew of cowpea, caused by Erysiphe polygoni, controlled with benomyl, 346;
---: Rhizoctonia sp., Sclerotium rolfsii,

and Pythium sp. controlled with potassium azide as seed treatment, 604
---: virus: mosaic virus causing mottling and blistering of foliage found in Puerto Rico, 212; high percentage of seedborne viruses and fungi found in cowpeas grown in eight States for seed purposes, 243; cowpea mosaic virus, vector efficiency of bean leaf beetle and cucumber spotted beetle greater with cowpea as test plant than with soybean, 1053
Crambe abyssinica: Alternaria circinans is cause of serious disease in Louisiana test plots to this potential new oilseed plant, 28
Cranberry: physiological breakdown in cranberries, and inhibition by intermittent warming during cold storage, 219
Criconemoides rusticum: 465
Cronartium comandrae: 471
--- fusiforme: 498
--- ribicola: 441; 1031
Cross-protection: no cross-protection observed between strains A and B of maize dwarf mosaic virus in grain sorghums, 627
Crown rot (see Rot)
Crownvetch (Coronilla varia): Sclerotinia trifoliorum infection, 811
Crucifers: cabbage: systemic infection of seedlings by Plasmodiophora brassicae race 6 found more frequently than previously reported, 453; control of clubroot with benomyl fungicide, 456; sugarbeet cyst nematode found on cabbage in Florida field, first report of occurrence in the State, 635
---: Crambe abyssinica (possible new oilseed plant) is host of Alternaria circinans in Louisiana, 28
---: radish, flecking and chlorosis of young plants from air pollution, 8
Cucurbits: cucumber: 18 cucumber varieties screened for resistance to target spot, downy mildew and powdery mildew, in Florida, 52; benomyl completely suppressed powdery mildew under greenhouse conditions, affected element accumulation, virus infection affected plant weight and element accumulation, but did not influence level of benomyl for mildew control, 141; nematicidal activity of 1,3 dichloropropene and 1,2-dichloropropane to three types of plant-parasitic nematodes, 196; rapid qualitative method with use of cucumber fruits to assay soil for Pythium and Phytophthora species, the primary cause of cucurbit root rot in Iran, 261
---: muskmelon: cucumber mosaic virus in cantaloup, 176; control of powdery mildew by root application of benomyl, in Israel, 232; control of Sclerotinia sclerotiorum on greenhouse-grown melons, 909
---: squash, air pollution damage in Cincinnati caused breakdown of chlorophyll between minor veins in leaves, 8
---: sugar pumpkin, nematode control with experimental nematicide, DuPont's 1410 nematicide-insecticide, 187
---: virus: cucumber is herbaceous host of cassava mosaic virus in India, 34; watermelon mosaic epiphytotic in New York in 1969 resulted in heavy losses among commercial species of Cucurbitaceae, 744; susceptibility of Cucurbita pepo and Citrullus lanatus introductions to watermelon mosaic virus-2, 880
---: watermelon: sporulation potential of Pseudoperonospora cubensis on watermelon, 108; powdery mildew of watermelon fruits reported from South Carolina for the first time, 372; control of Fusarium wilt with benomyl soil drench, 561
Culture: monoxenic culture of Ditylenchus dipsaci and Pratylenchus penetrans on onion and alfalfa callus produced on modified Krusberg's and White's media, 251; wheat grown in mist culture under water stress and inoculated with stripe rust, showed greater reduction in root growth than with either condition alone, 387; cultural, biochemical and serological tests for determination of Pseudomonas syringae as pathogen of Austrian winter pea, 467
Cyamopsis tetragonoloba (guar): susceptibility to new virulent race of Xanthomonas cyamopsidis, 111
Cylindrocladium crotalariae: 320
--- floridanum: 320
--- scoparium: 320
Cylindrosporium ulmicola: first record of pathogen in Argentina, 939
Cytospora sp.: 179; 996
--- spp.: 832

Damping off: of crimson clover, damping-off fungi controlled with 5-minute exposure to radio-frequency energy (rf) to soil, 268
---: bean: of snap bean, additional sources of resistance to disease found in several dark seeded plant introduction accessions, 40; of mungbean, in Iran, 246
---: of cotton seedlings, Rhizoctonia solani controlled with Vitavax, 524
---: of pea seedlings: due to Pythium sp., 17 of 22 formulations, reduced loss but thiabendazole combined with other fungicides excelled all other treatments in seedling production, 349; Rhizoctonia

solani controlled with benomyl, 1049
Date palm (Phoenix dactylifera): Graphiola leaf spot reduced chlorophylls a and b in moderately and severely infected varieties of date palm in India, 617
Datura spp.: classification of 37 virus isolates from arborescent Datura species into five groups, Datura species may act as reservoirs for viruses that infect solanaceous species, 675
Decline: of live oak, research on this disease complex indicates that wilting symptoms are caused by mechanical interference with water flow, the presence of a toxin, or both, 669; of lucerne stands in Western Australia, Fusarium oxysporum pathogenic to seedling roots, 1061
Desmodium canum: Desmodium mosaic virus isolated from D. canum appears to be member of potato virus Y group, 161
Dew: 8 hours minimum of dew, temperature above 10°C, followed by 24 hours above 10° favors establishment of stem rust infection on wheat, 959
Dew recorder: a new approach to recording the wetting parameter by use of electrical resistance sensors, 474; modified version of the Taylor dew meter that will operate automatically for 8-day periods, 512
Diabrotica undecimpunctata howardi: 1053
Diagnostic symptoms: of pea fizzletop virus, 355
Diaphorina citri: 1092
Diaporthe alleghaniensis: 16
Dibotryon morbosum: 217
Dieback: twig dieback of Abies spp., perithecia of a Pleospora sp. developed on primary cankers of dead, curled twigs, 700; of 'Robinson' tangerine, benomyl spray in March more effective than in June, 869
Differential hosts: for strains A and B of maize dwarf mosaic virus, 630
Differential reaction: of Pythium sp. and five isolates of Rhizoctonia solani to fungicides on pea seedlings, 349
Digestibility: of Johnson grass and a sorghum-Sudan grass hybrid infected with maize dwarf mosaic virus, 424
Diphtherophora communis: first report of this nematode species in Iran, 339
Diplodia gossypina: 29; 586; 796
--- maydis: 658
--- zeae: 624
Dips: fruit dips for control of postharvest decay of Anjou pears, 887; for gladiolus corms, benomyl and thiabendazole reduced corm rot, 689
Diseases: fungus and bacterial diseases of plants in Cordoba, Argentina, 939
Distribution: of sorghum downy mildew (Sclerospora sorghi) in the U.S., 311; varietal and geographical distribution of walnut phloem canker and bark canker in California, 373; of Dutch elm disease in Minnesota, 706; of Dutch elm disease in the United States in 1970, 929; origin and spread of southern corn leaf blight in the U.S. in 1970, 1104
Ditylenchus dipsaci: 251
Dogwood: control of Meloidogyne incognita in seedlings by chemical dips, 952
Dollar spot: Sclerotinia dollar spot, of turfgrass, excellent control with benomyl and thiabendazole, 1080
Dothiorella quercina: 669
Dothistroma pini: 403; 679
Douglas-fir (see Pseudotsuga)
Downy mildew: cucurbits: cucumber varieties Poinsett and Cherokee showed resistence to causal organism, Pseudoperonospora cubensis, 52; sporulation potential of P. cubensis on two watermelon cultivars measured, 108
---: of sorghum: Sclerospora sorghi now found in 10 States, control measures urgently needed, 311; inoculation of seedlings by use of infected leaves placed in direct contact with sorghum seedlings, infection occurred readily through stomatal openings, 603
---: of soybean, immunity to Peronospora manshurica, 901
Downy spot: of pecan, pre-pollination applications of benomyl, dodine, and triphenyltin hydroxide were best of several tested for control, 725
Dutch elm disease: Ceratocystis ulmi, inoculated into base of 12-foot tree, colonized half the main stem within 3 days and all of the main stem in 6 days, 58; symptom progression in elm following inoculation, 127; trunk injections and application to soil with thiabendazole did not give control in Ohio, 240; distribution in Colorado in 1969, 266; identified in Wyoming, 547; slow spread in Idaho, 701; the present known distribution, pattern of spread and distribution of the smaller European bark beetle in Minnesota, 706; distribution in 1970 in the United States, 929; detection in Texas for the first time, 931; attempts to control disease by pruning are neither effective nor practical, 985
Dwarf mistletoe (see Mistletoe)

Ear cockle: of wheat, in India caused by nematode, Anguina tritici, 594
Ecuador: 675
Eelworm (see Nematodes)
Eggplant (Solanum melongena)(brinjal):

Verticillium wilt: found for the first time on eggplant in Turkey, previously recorded only on cotton, 241; grafting eggplants on resistant tomato rootstocks gave eggplants of desirable type with high level of resistance to wilt, 846
---: virus: mosaic disease on 'Pusa purple Long', vectors, host range and physical properties of virus determined, 437
Electrical resistance sensors: for measuring dew or wetness, instrumentation differs in three major ways from previously described systems, 474
Element accumulation: benomyl apparently affects accumulation of certain elements in cucumber plants, depending upon dosage, 141
Elevation: elevation and temperature effects on severity of maize dwarf mosaic virus in sorghum in Arizona, 1064
Elm: Dutch elm disease: 58; 127; 240; 266; 547; 701; 706; 929; 931; 985
---: Siberian, flux from wetwood-affected trees, caused by Erwinia nimipressuralis, drained by means of plastic tubing, 66
Endothia parasitica: 713
England: 150; 370; 734; 802
Environmental factors: severity of infection by Fusarium roseum of 'Merion' Kentucky bluegrass correlated with soil temperature and host nutrition, 1018
Epidemiology: bacterial canker, of French prune, 332; of southern corn leaf blight, exploratory experiments, 1131
Epiphytotic: peanut stunt virus caused a destructive disease of beans in North Carolina, 183; of red rot of sugarcane in Burma in 1969, 427; of watermelon mosaic among Cucurbitaceae in central New York in 1969, 744; of southern corn leaf blight in the United States, 1099-1136
Erwinia amylovora: 96; 203; 576; 638; 835; 1039
--- atroseptica: 315
--- carotovora: 263; 863; 964
--- nigrifluens: 373
--- nimipressuralis: 66
--- rubrifaciens: 373
Erysiphe cichoracearum: 52; 372; 399
--- graminis: 91
--- polygoni: 346; 399; 606
Evaporation: effect of evaporation rate as indicator for potato late blight development, 768

False broomrape: on sweetpotato, 979
Fern: leatherleaf (Polystichum adiantiforme): control of Pratylenchus penetrans with nematicides, 1016
Fertilizer: high rates of nitrogenous fertilizers applied to rice plants increased susceptibility to blast disease, 632
Ficus elastica (rubber plant): pathogenicity of Cercospora sp. to ornamental rubber plant in Florida, 199
Field spread: of barley stripe mosaic virus in Montana, 749
Fiji: 1035
Filbert (Corylus avellana): filbert stunt proposed as name for undetermined disease of filbert, in Oregon, 69
Fire blight (see Blight)
Fizzletop disease: of pea, virus disease of pea that affects ovule development, 355
Flat apple (see Apple virus)
Florida: 52; 173; 199; 287; 411; 569; 608; 635; 678; 679; 759; 760; 827; 865; 869; 1127
Flower blast: of statice, one of three diseases limiting economic importance, in Florida, 759
Fluorescence: internal fluorescence of cotton plants from Arizona found unrelated to fiber fluorescence associated with Aspergillus flavus boll rot, 408
Foam: urethane foam useful in tree cavity fills, 858
Fomes annosus: 330; 438; 872; biological control with oidia of Peniophora gigantea applied to stumps, 799
Forecasting: plans for forecasting corn blight epidemics, 1134
Fragaria ananassa (see Strawberry)
France: 915
Freckle scurf: bark disorder of 'Winesap' apple, 44
Fruit blotch: of apple, transmissible and non-transmissible fruit blotching syndromes tested in British Columbia, 490
Fruit distortion: distortion ringspot virus caused abnormalities in fruit formation of papaya cultivars, 181
Fungi: presence of internally-borne fungi in southern pea seed is influenced by location of seed production, 243; control of Pythium sp., Rhizoctonia sp., and Sclerotium rolfsii, with potassium azide seed soak on cowpea, 604; Pythiaceous fungi inhibited by nematicide DBCP, in cultures and in the field, 622; species of fungi associated with cotton boll rot in the Yazoo-Mississippi Delta, 1966-1968, 796
Fungicides: 2 of 11 experimental fungicides eliminated Ustilago nuda from leaves of 'Larker' barley, 927
---: benomyl: reduction of tomato Fusarium wilt symptoms appears to depend on an effective level of fungitoxicant while fungus is spreading through plant, 12; as spring application was more effective than thiabendazole or urea in reduction

of ascospore discharge of Venturia inaequalis from overwintered apple leaves, 27; controlled damping-off of cotton seedlings, 76; effective in control of apple powdery mildew and cherry leaf spot in Poland, 81; postharvest dip more effective than preharvest spray for control of Nectria decay of apple, 83; controlled powdery mildew of wheat at Prince Edward Island, Canada, 91; benomyl better than thiabendazole under commercial conditions, for control of stem-end rot of banana fruit, 93; benomyl and DMTT effective in suppressing bean root rot from Thielaviopsis basicola, 114; controlled root rot of potted tobacco, apparently by eradication of T. basicola, 136; virus infection had little effect upon systemic control of powdery mildew of cucumber by benomyl, element accumulation changes varied with the dosage of the fungicide, 141; as dormant spray in October or March, controlled Ustilago striiformis on 'Merion' Kentucky bluegrass in Pennsylvania, not effective in period of active grass growth, 146; best os several for control of stripe smut on 'Merion' Kentucky bluegrass; it did not control Helminthosporium leaf spot, 168; soil drench plus Tween 20 appears to reduce rate of symptom development of Verticillium wilt in potato plants, 171; root application to control powdery mildew of muskmelon in Israel, 232; benomyl and thiabendazole as soak or dust effective on iris bulbs infected with Fusarium basal rot, 235; best of four fungicides against Verticillium wilt in artificially infested potato seed tubers, 254; benomyl and thiram compared for control of Fusarium rot of gladiolus, 289; benomyl and thiabendazole superior to SOPP and 2-aminobutane (2-AB) in control of Penicillium digitatum on three orange varieties, 325; as soil drench on rooted cuttings and plants of poinsettia, controlled Thielaviopsis root rot, 328; used as soil drench, seed treatment or foliar spray, controlled powdery mildew of cowpea, 346; in greenhouse tests, foliar sprays applied at 3-week intervals effectively controlled powdery mildew of strawberry, 362; benomyl and thiabendazole for control of Fusarium basal rot of Narcissus, 377; benomyl and thiabendazole mixed with a rooting hormone increased rooting percentage of rhododendron cuttings, 427; prevented cabbage root hair infection by Plasmodiophora brassicae in transplants, 456; for control of Septoria leafspot of celery, 531; controlled Fusarium wilt of potted tomato and watermelon, in Greece, 561; very effective against powdery

Fungicides: benomyl (continued):
mildew and Botrytis blight of hydrangea, 606; benomyl and Cercobin controlled Pezicula spp. infection on apple, and were superior to mercurial formulations, 640; as preharvest spray on 'D' Anjou' pear fruit controlled postharvest decay, 647; benomyl and thiabendazole as preharvest dips greatly reduced blue mold decay of pears, 655; reduced Pratylenchus penetrans over 50% and reduced severity of weather fleck of tobacco until late harvest, in Connecticut, 672; benomyl + polyvinyl chloride better than benomyl alone for control of Dothistroma needle blight of Austrian pine, 679; benomyl and thiabendazole effective against Fusarium corm rot of gladiolus, 689; superior control against Botrytis cinerea, cause of raceme blight of macadamia, 694; benomyl, dodine and triphenyltin hydroxide were best of several for control of downy spot of pecan, 725; surfactant PE-510 most effective of several tested as an adjuvant for benomyl to prevent Verticillium symptoms after stem inoculation of pot-grown cotton plants, 741; control of postharvest decay of peaches, 764; as a preventive spray in spring is effective against Fusarium blight of 'Merion' Kentucky bluegrass, 771; after use on bentgrass turf in Australia, a patch disease developed, caused by a Basidiomycete, 774; best of fungicides tested to reduce numbers of perithecia of citrus greasy spot fungus, 865; reduced incidence of dieback in 'Robinson' tangerine, 869; best of several tested for control of Botrytis fruit rot and powdery mildew of strawberry, 883; benomyl and thiabendazole dips gave good control of decay in cold-stored Anjou pears, 887; benomyl, cycloheximide and Difolatan controlled mouldy rot of tapping panel of Hevea in Malaysia, 897; controlled Sclerotinia sclerotiorum on muskmelons in Israel, but a benomyl-tolerant strain of fungus was observed, 909; controlled Cercospora leafspot of peanut in Virginia tests, 955; as soil amendment, 981; seed dressing or soaking, or soil drenching controlled Rhizoctonia damping-off of peas, 1049; gave excellent control of Sclerotinia dollar spot, good control of red thread and Rhizoctonia brown patch but no control of Helminthosporium leafspot of turfgrass in Pennsylvania, 1080

---: BSM-11 (50% potassium 2,4,6-trichlorophenate plus 10% phenylmercuric

acetate), eliminate the stem canker fungus of blueberry, in dormant cutting wood, 122
Fungicides: Bordeaux, effective for control of Dothistroma needle blight of Austrian pine, 679
---: captan, over a 15-year period has reduced rotten strawberry fruit by about one-third, and increased yield and net profit in Washington, 132; as postharvest dip for strawberries was inferior to several treatments, 431; captan, dithianon and ziram most effective fungicides tested for control of peach brown rot, in New South Wales, 776
---: chloroneb: controlled Typhula blight of bentgrass in Michigan, 1075; chloroneb and PCNB gave no protection for stripe smut of bluegrass, but good control against Helminthosporium leaf spot, 168
---: copper: basic copper sulfate or ferbam at petal fall and summer application of zineb-oil will control Gloeosporium foliicolum, fungus disease new in Florida, 173; copper salts of fatty and rosin acids (TC-90), effective in control of Dothistroma pini on Austrian pine, 679
---: cycloheximide: cycloheximide dip eradicated powdery mildew of rose seedlings, 598
---: DCNA (2,6-dichloro-4-nitroaniline) used as dip, reduced spoilage from Rhizopus stolonifer if storage temperatures were 60°F or above, 486; fungicide dip for reduction of postharvest decay of peaches, 764; DCNA + captan and DCNA + Difolatan evaluated as seed protectants on 15 crops at several locations, 1041
---: dichlone and phenylmercury nitrate, added to foam for tree cavity fill reduced mycelial growth of white- and brown-rot fungi, 858
---: Difolatan, at 125 ppm gave excellent control of Botrytis cinerea on macadamia under bioassay, and under field conditions, 694
---: dodine: dodine-captan spray injury, on peach and nectarine, captan is apparently the toxic agent, 152; dodine, for downy spot of pecan, 725
---: HOE 2873, gave good control of powdery mildew of hydrangea but no control of Botrytis blight, 606
---: maneb: maneb, ferbam and thiram, in the laboratory, showed the greatest inhibition of blister canker of pear, 296; maneb and ziram controlled Scleroderris canker on red pine nursery stock in Wisconsin, 663; effective in control of needlecast, caused by Lophodermium pinastri, on red and Scotch pines, 731; maneb-captan, for control of Cerospora leafspot of statice, 760; coordination of zinc ion and maneb controlled anthracnose of oriental poppy, 1037
---: O-199 (O,O-diethyl phthalimidophosphonothioate), effective against powdery mildew of rose seedlings when used as a dip, 598
---: organo mercurials, for seed treatment of oats, increased tolerance by Pyrenophora avenae observed in New York State, 783
---: oxytetracycline, weekly sprays necessary for effective control of peach bacterial blight in Maryland, 208
---: potassium azide (KN_3), fungicidal activity of herbicide shown in seed treatment with 'Early Ramshorn' cowpea, 604
---: streptomycin with sulfur and glyodin as a night spray for control of fire blight of apple, 203
---: tetrachloroisophthalonitrile (chlorothalonil), for statice, 760
---: thiabendazole, reduced ascospore discharge of apple scab, 27; not effective as a control for Dutch elm disease, by soil application or by injection into trees, 240; in combination with three other fungicides reduced seed decay of pea more than any of the other 18 formulations tested, 349; controlled septoria leafspot on celery, 531; controlled blue mold decay of pear, 655; effective against corm rot of gladiolus, 689; control of postharvest decay of peaches, 764; as amendment, 981; gave fine control of Sclerotinia dollar spot of turfgrass, but little control of red thread, Rhizoctonia brown patch or Helminthosporium leafspot, 1080
---: thiram: thiram when coupled with an insecticide controlled onion smut and onion maggot at 0.5-, 1.0-, or 1.5-inch planting depths, 290; effective as seed treatment to control Helminthosporium spiciferum on Bermuda grass, 358; thiram-captan drench effective against Rhizoctonia stem rot of chrysanthemum, 679; thiram + 10% 2-chloro-4-(hydroxymercuri)phenol, for control of Fusarium blight of 'Merion' Kentucky bluegrass, 771
---: triphenyltin hydroxide, for downy spot of pecan, 725
---: urea, to control discharge of ascospores of Venturia inaequalis, 27
---: Vitavax (5,6-dihydro-2-methyl-1,4-oxathiin-3-carboxanilide) showed systemic action against Rhizoctonia solani on cotton seedlings, 524
---: zineb: weekly spray on corn varieties

reduced infection with Puccinia sorghi and increased yields in India, 788; controlled Cercospora leafspot of statice, 760
---: ziram, effective against Scleroderris canker of red pine, 663; preharvest spray for control of postharvest decay of 'D' Anjou' pears, 647
---: several tested together: 22 chemicals tested for systemic fungitoxic activity against Sclerotium rolfsii and Thielaviopsis basicola, 223; to reduce damping-off of pea seedlings from Rhizoctonia solani and Pythium sp., 349; five effective against Septoria apiicola on celery in California with 14-day preventive spray schedule, 531; for control of raceme blight of macadamia, caused by Botrytis cinerea, in Hawaii, 694; several fungicides in combination with nematicides increased yield of strawberries in West Virginia, 923; Difolatan, Polyram, or maneb combined with insecticide EPN greatly reduced spread of cotton boll black rot in Taiwan, 29; approved chemicals as protectant fungicides against Helminthosporium maydis, 1112
Fusarium sp.: 28
--- spp.: 268; 586; 796
--- moniliforme: 307; 483
--- oxysporum: 1061
--- --- f. sp. gladioli: 235; 289; 689
--- --- f. sp. lycopersici: 12; 561
--- --- f. sp. narcissi: 377
--- --- f. sp. niveum: 561
--- roseum: 771; 1018
--- --- f. sp. dianthi: 446
--- solani f. phaseoli: 698
--- vasinfectum: 324
Fusicladium effusum: 829

Gardenia jasminoides: nematicides as soil drenches, or granules worked into soil gave best control of Meloidogyne incognita, 966
Genetic control: susceptibility of corn inbreds and crosses with Texas male-sterile cytoplasm to Phyllosticta sp., suggests genetic factors are involved, 277; 281
Georgia: 12; 60; 419; 465; 586; 620; 952; 987; 1039; 1077
Geotrichum candidum: 881
Geranium (see Pelargonium)
Gibberella zeae: 658
Gladiolus: surveys of 1968 and 1969 indicate that 20.9 and 27.2% of all field-grown gladioli in North Carolina showed virus symptoms, 125; Fusarium rot and bacterial scab, corm treatment with benomyl and thiram compared for control, 289; Fusarium corm rot, benomyl and thiabendazole both reduced corm rot, 689
Gloeosporium cingulata: 1037
--- foliicolum (Glomerella cingulata): 173
--- musarum: 93
Glomerella gossypii: 586
Gnomonia caryae: 408
--- --- var. pecanae: 408
Grafting: of eggplant on to resistant rootstocks to control Verticillium wilt, 846
Granadilla (Passiflora quadrangularis): fruit rot, caused by Botryodiplodia theobromae, 809
Grape (Vitis vinifera): identification of Sclerotinia laxa on grape in New Zealand is apparently a new record of distribution, 504; (Vitis rupestris) whole plant cured of grapevine fanleaf by thermotherapy, 915; incidence of viruses in old abandoned California vineyard found to be low and fanleaf was completely absent, 1055
Graphiola phoenicis: 617
Grasses: creeping bentgrass: Pyrenochaeta terrestris is root pathogen of bentgrass in Illinois, 506; benomyl apparently predisposed bentgrass turf to a patch disease caused by an unidentified Basidiomycete, 774; chloroneb effective in control of Typhula blight, 1075
---: Bermuda grass, seed infested with Helminthosporium spiciferum, apparently a first report of occurrence, thiram as seed treatment gives excellent control, 358
---: 'Merion' Kentucky bluegrass: 6 oz or more of benomyl applied in October or March in dormant period suppressed development of stripe smut, 146; fall application of benomyl effectively controls stripe smut but gives no protection against Helminthosporium leaf spot, 168; Kentucky bluegrass, symptomatology of stripe-smutted inflorescences, in Iowa, 206; chemical control of Fusarium blight, best with spring preventive spray of benomyl, 771; high soil temperature, high nitrogen levels or low moisture predispose grass to infection by Fusarium roseum, 1018
---: Johnson grass and sorghum-Sudan grass hybrid, maize dwarf mosaic virus infection caused some reduction in yield in sorghum-Sudan grass hybrid, not in Johnson grass and did not affect protein content or digestibility of either grass, 424
---: Panicum repens, a common weed in rice paddies identified as a collateral host of Cercospora oryzae, 272

---: turfgrasses, soil assays in South and Southeast from problem golf greens revealed many Pythium species present, 419; Meloidogyne naasi is parasitic to several species in California, 940; Rhizoctonia solani and Sclerotinia homoeocarpa inhibited with benomyl and thiabendazole, 981; benomyl and thiabendazole compared to standard fungicides such as maneb and zineb for control of turfgrass diseases, 1080
Grassy stunt disease: of rice, development of mass screening method for testing resistance, 565
Gray rot: of lettuce, 964
Graywall: of tomato, in Florida, tobacco mosaic virus is suggested as causal agent, 678
Greece: 73; 134; 561
Greening disease: of citrus, in Nepal, 1092
Green mold: of oranges, caused by Penicillium digitatum, benzimidazoles effective in control, 325
Green mottle virus (see Apple virus)
Groundnut (see Peanut)
Guar (Cyamopsis tetragonoloba): new virulent race of Xanthomonas cyamopsidis pathogenic to three formerly bacterial blight resistant cultivars of guar, in Texas, 111

Hackberry (Celtis laevigata): Polyporus zonalis associated with stem breakage of hackberry in Alabama, 322
Hawaii: 694
Helicotylenchus nannus: 339
--- pumulis: 752
Helminthosporium sp. 1080
--- maydis (Cochliobolus heterostrophus): 501; 708; 819; 1099-1136
--- papulosum f. pyri: 296
--- sorokinianum: 594
--- spiciferum: 358; 405
--- turcicum: 896
--- vagans: 168
Hemileia vastatrix: 355; coffee rust now firmly established on coffee in Brazil, 539
Hemlock (see Tsuga)
Heterodera avenae: 134; 594
--- cacti: 692
--- glycines: 544
--- rostochiensis: 261; 306
--- schachtii: 196; 635
--- tabacum: 25; 781
Hevea brasiliensis (Rubber): fungicides benomyl, cycloheximide and Difolatan controlled Ceratocystis fimbriata, cause of mouldy rot of the tapping panel, 897
Hibiscus cannabinus (see Kenaf)
--- sabdariffa (see Roselle)
Highway plantings: salt damage to eastern white pine from wintersalting of highway, 1021
Hoplolaimus galeatus: 752
Hormone: indole butyric acid, a hormone powder mixed with benomyl and thiabendazole significantly increased rooting of rhododendron cuttings over fungicide alone or hormone alone, 427
Host effects: of pea from virus-induced fizzletop disease, 355
Host list: of previously unreported powdery mildews found in California, 399
Host range: of bromegrass mosaic virus, 191; of cowpea mosaic virus, 212; of cherry short stem virus from Oregon, 651; for Meloidogyne naasi, 940
Hot water treatments (see Temperature)
Hungary: 286
Hydrangea: suppression of Erysiphe polygoni and Botrytis cinerea with benomyl, 606
Hylemya antiqua (onion maggot): 291
Hyperparasitism: in true mistletoes (Phoradendron spp.) in California, and also autoparasitism, 15
Hypoxylon atropunctatum: 669
--- pruinatum: 462

Idaho: 497; 701
Illinois: 49; 67; 422; 501; 506; 832; 973; 1095
India: 34; 181; 226; 272; 337; 391; 437; 453; 594; 617; 727; 788; 976; 1049
Indiana: 583
Infectivity dilution curves: of soilborne wheat streak mosaic virus, 550
Infraspecific forms: of soybean cyst nematode and identity of four "races", 544
Inoculation: methods tested for inoculation of strawberry with red stele pathogen, 739; studies with Fusicladium effusum using excised pecans, 829
Inoculum: conidial infection with Sclerospora sorghi obtained in sorghum seedlings by using natural deposition of conidia on seedling leaves from infected corn and sorghum plants, 603
Insect collection: collection of large quantities of insects for experimental use with a compressed air aspirator, 56
Introductions: tomato accessions tested for anthracnose resistance, 32
In vitro: inhibition of Verticillium alboatrum on agar by benomyl using six different surfactants as adjuvants, 741
Iowa: 206; 624
Iran: 246; 261; 339; 1085
Ireland: 83
Iris: benomyl and thiabendazole were usually more effective and less phytotoxic than mercurials for treatment of Fusarium-infected iris bulbs, 235
Irrigation ponds: Pythium myriotylum from

irrigation ponds in Georgia found to be pathogenic to tomato, snapdragon, pansy and sweet william, 1077
Isolates: cucumber mosaic virus isolates from sugarbeet, lettuce and cantaloup exhibited only minor variation in incubation period and disease severity within 14 selected hosts, 176; variation in telial formation in certain isolates of Puccinia penniseti, 646; of Sclerotinia trifoliorum, showed no specificity for either alfalfa or crown-vetch, 811; Illinois isolate of Meloidogyne naasi pathogenic to 'Wintok' oats, 972; of Corynebacterium michiganense, variation in virulence, 976
Isolation: a portable tubular framed chamber designed for containment of infected plants and exclusion of airborne fungal contaminants, 130; isolation, inoculation, temperature relations and culture of a Cercospora pathogenic to Ficus elastica 'Decora', 199
Israel: 232; 325; 442; 643; 735; 768; 804; 909
Italy: 3; 389

Kenaf (Hibiscus sabdariffa): Meloidogyne incognita acrita pathogenic to all 28 varieties tested in Nigeria, 547
Kentucky: 723

Laboratory: temporary quarantine facility for research on non-resident pathogens with safeguards against their escape, 3
Lake States: 731
Laurel (Kalmia latifolia): Fomes annosus infection of mountain laurel in North Carolina is apparently a new host record, 560
Leaf blotch: of oat, possible development of a mercury-tolerant strain of Pyrenophora avenae in New York, 783
Leaf chlorosis: of pear, symptoms caused by Alternaria tenuis in greenhouse inoculation identical to those observed under field conditions, 871
Leaf spot: of celery, caused by Septoria apiicola, controlled with several fungicides with 14-day spray schedule in California, 531; of cherry, caused by Coccomyces hiemalis, controlled with benomyl in Poland, 81; Graphiola leaf spot of date palm, caused by Graphiola phoenicis, 617; of moth bean (Phaseolus aconitifolius) in India, 976; of mung bean (Phaseolus aureus), caused by Alternaria alternata, in India, 453; Cercospora oryzae on rice, a new collateral host, Panicum repens, has been found in India, 272; of statice, Cercospora leafspot, 759; several fungicides effective in control on statice, 760; of tobacco, caused by Cercospora nicotianae, in United Arab Republic, 666; of wheat, caused by Helminthosporium sorokinianum, 594
--- and fruit spot: of sweet pepper, caused by Corynebacterium michiganense, in Israel, 804
Leatherleaf fern: root lesion nematodes, Pratylenchus penetrans populations reduced on fern transplant material with six nematicides, 287
Lebanon: 435
Lettuce: corky root rot, resistance in lettuce, 754
---: market diseases: New York market: bacterial soft rot and gray mold in western lettuce comprised 90% of all decays, 263; brown stain of western head lettuce, 856; diseases of head lettuce from California, 964
---: nematodes: root-knot nematodes, three nematicides effective in reduction of nematode populations in New York, 368
Lily (Lilium longiflorum): aldicarb, carbofuran and methomyl nematicides as direct applications to bulbs significantly increased bulb weight over accepted phorate treatment, 1012
Liriodendron tulipifera (Yellow-poplar): three Cylindrocladium species pathogenic to potted seedlings, 320
Lirula nervisequia conspicua: 275
Little leaf: of apple, a probable temperature-induced genetic disorder of apple seedlings, 1059
Liver spot disease: of pecan, caused by Gnomonia caryae var. pecanae, reported for the first time in Alabama, 408
Longevity (see also Survival): longevity and pathogenicity of several isolates of Stemphylium species stored under mineral oil, 580
Lophodermium pinastri: 731
Louisiana: 28; 554; 713; 815
Lucerne: a decline condition, caused by Fusarium oxysporum, 1061

Macadamia: raceme blight of macadamia, caused by Botrytis cinerea, benomyl and Difolatan gave superior control in laboratory bioassy and in field control, 694
Macrophoma sp.: 315
Macrophomina phaseoli: 286; 891; 1088
Maize(see Corn)
Malaysia: 897
Male-sterile cytoplasm: its possible role in susceptibility of inbred lines of corn to Phyllosticta sp., 277; corn hybrids or inbreds containing Tms showed increased susceptibility to Phyllosticta

leaf blight but all were not equally susceptible, 281; increased susceptibility of corn inbreds and hybrids with Tms cytoplasm to Helminthosporium maydis indicates a maternal inheritance of susceptibility, 501; reaction of corn seedlings with male-sterile cytoplasm to H. maydis, 708; in susceptibility to southern corn leaf blight, 1099-1136
Mango (Mangifera indica): Verticillium wilt found in young trees in Florida planted where tomatoes had previously grown, 569
Market diseases: lettuce: gray mold rot (Botrytis cinerea) and bacterial soft rot (Pseudomonas sp. and Erwinia carotovora) of lettuce on New York market caused 90% of all decays, 263; brown stain of head lettuce, shipped from California, 856; diseases of head lettuce from California, 964
---: stem-end rot of Chinese gooseberry (kiwi fruit), Phomopsis sp. recovered, 422
Mayland: 208; 891; 1059
Massachusetts: 296
Meloidogyne spp.: 324, 411
--- incognita: 952; 966; 1014; 1056
--- --- acrita: 196
--- naasi: 940; 973
Methods and techniques: bacterial canker resistance determinations of seedlings obtained without destroying them, 36; simple technique for evaluating translocation of fungicides, 223; control of seedling damping-off of crimson clover in greenhouse soils by radio frequency energy, 268; seedling production of peas treated with fungicides was greater in paper towels than in plastic boxes, 349; incubation of stump samples on wire racks in large containers effective for detection of Fomes annosus infection, 438; use of insect vector, Nilaparvata lugens for mass screening of rice varieties for resistance to grassy stunt, 565; sterile mineral oil for long-term preservation of Stemphylium species, 580; use of detached leaves for screening tomato plants for single gene resistance to late blight race O, 681; evaluation of fungicides by laboratory bioassay for control of Botrytis cinerea on macadamia in Hawaii, 694; adjuvant effects of surfactants on benomyl inhibition of Verticillium on Czapek solution agar, 741; scoring system for rating severity of fire blight in pear, 835; rapid technique for making leaf-tissue grafts to transmit citrus viruses, 907; elimination of grapevine fanleaf in grape cv. du Lot by thermotherapy, 915; an open-ended system for differentiating races of stripe rust of wheat in the United States, 992; breeding methods to produce hybrid corn seed, by use of normal cytoplasm, male-sterile cytoplasm or a blend of normal and male-sterile cytoplasms, 1118; forecasting corn blight diseases, caused by Helminthosporium maydis and other Helminthosporium species, 1134
---: equipment: temporary quarantine laboratory for research on imported plant pathogens, in Rome, Italy, 3; remote sensing and recording of relative humidity by electric psychrometer; a computer program is also available, 19; a portable chamber for the growth and isolation of infected plants, 130; a modified roller type mill for treating seeds, 414; a dew recorder that yields a very accurate record of wetting duration as well as an estimation of degree of wetness with electrical resistance sensors but which differs in three major ways from previously used grids and the electrical resistance technique, 474; a new spore sampler makes discrete spore prints for specified time intervals, 498; an 8-day dew recorder, a further modification of the Taylor meter, 512; static cage technique for transmission of cotton leaf curl virus by single insect vectors, 528; an improved method for concentrating nematodes from the Baermann apparatus, 899
---: inoculation: compressed-air insect aspirator useful for collection of large numbers of insects for inoculation experiments, 56; three inoculation techniques evaluated for determination of fire blight resistance in pear seedlings, 96; artist's airbrush inoculation of soil-borne wheat mosaic virus increased percentages of infection over the manual leaf rubbing method, 550; use of conidia from infected corn and sorghum leaves to inoculate sorghum seedlings, 603; inoculation of corn ears by spraying with spore suspension of Gibberella zeae and insertion of toothpicks overgrown with G. zeae into ear shoots, 658; use of macerated mycelium of Phytophthora fragariae effective for inoculation of strawberry roots, 667; development of a rapid, satisfactory innoculation technique to evaluate large numbers of strawberry plants for resistance to Verticillium albo-atrum, 682; comparison of four methods of inoculation of strawberry with Phytophthora fragariae, 739; simultaneous inoculation of pine stumps with oidia-oil of Peniophora gigantea in the power saw suppressed Fomes annosus, 799; use of excised pecans for inoculation studies with Fusicladium effusum, 829; spe-

cialized inoculation technique used to show variation in virulence in Corynebacterium michiganense isolates, 976
---: isolates: methods for isolating a Cercospora sp., and proof of its pathogenicity to Ficus elastica, 199; use of cucumber fruit as a simple method to isolate Phytophthora and Pythium species from soil, 261;purification technique and production of telia in certain isolates of Puccinia penniseti, 646
Mexico: 762
Michigan: 16; 290; 612; 635; 849; 853; 985; 1075
Migration: migration and penetration of Erwinia amylovora in artificially inoculated apple tree tissue, 576
Mineral oil: as a long-term preservative for Stemphylium species, 580
Minnesota: 132; 460; 706
Mississippi: 796
Missouri: 203; 319
Mist culture: stripe rust and water stress together reduced root growth of wheat more than either factor alone, 387
Mistletoe (Phoradendron spp.): Phoradendron villosum subsp. villosum found parasitizing P. bolleanum subp. pauciflorum in California on Abies concolor; also P. juniperinum subsp. libocedri was found parasitizing P. bolleanum subsp. pauciflorum in same tree, 15
---: dwarf: of limber pine, now found 90 miles farther east in Montana, 441; two additional dwarf mistletoes found on the rare Brewer spruce, in Oregon and California, 488
Modified atmosphere: for control of postharvest decay of California strawberries, 431
Moisture: high soil moisture levels critical for infection of black walnut seedlings by Phytophthora citricola, 583
Molya: of wheat, caused by Heterodera avenae, of major importance in west India, 594
Monilinia fructicola: 764; 776
Monilochaetes infuscans: 1056
Montana: 441; 749
Moth bean (Phaseolus aconitifolius): new leaf spot disease, caused by Colletotrichum truncatum identified in India, 976
Mouldy rot: of Hevea brasiliensis, caused by Ceratocystis fimbriata controlled with fungicides, 897
Multiple sprouting disease: of citrus, a virus-like disease was transmitted mechanically from leaves and flower petals to Chenopodium quinoa, 1003
Mungbean (Phaseolus aureus): Rhizoctonia stem canker reduced plant populations in Iran by 4-57%, none of 40 types tested showed resistance, 246; leaf spot, caused by Alternaria alternata, apparently a first record of pathogen on mung, 453
Musa spp.: abaca and banana plants with symptoms of vascular wilt found in Philippines, Pseudomonas solanacearum apparently involved in wilt problem of abaca, but not primary agent for banana, 516
Mycoplasma: safflower phyllody, on basis of electron microscopy the etiologic agent of safflower phyllody is considered to belong to leafhopper-borne mycoplasmas, 735; mycoplasma-like bodies and viruses in Opuntia tuna with witches' -broom disease, 851; suppression of Philippine greening disease of citrus with antibiotics suggests etiologic agent is a mycoplasma-like organism rather than a virus, 1007
Mycosphaerella sp.: 865
--- caryigena: 725
Myzus persicae: 370; 437; 876

Narcissus: Fusarium basal rot control with benzimidazole fungicides, 377
Nebraska: 557; 572
Necrosis: leaf, of spinach, new disorder in Texas thought to be physiogenic in nature, 155
---: root, of strawberry, 8 species of Pythium isolated from diseased roots, 49
Nectarine: peach canker, caused by Cytospora spp., 832
Nectria galligena: 83
Needle blight (see Blight)
Needle cast: of bristlecone fir, caused by Lirula nervisequia var. conspicua, this is a first report of a new tree disease in the United States and the first known report of a disease on bristlecone fir, 275; Swiss needle cast of Douglas-fir, first report from the Lake States, 612; Lophodermium pinastri caused severe damage to several million Pinus resinosa and P. sylvestris seedlings in the Lake States, 731
Needle injury: of ponderosa pine, caused by air pollutants, 945
Nematicides(see also Antibiotics, Chemicals, and Fungicides): aldicarb, carbofuran and methomyl applied to lily bulbs increased bulb weight more than phorate treatment, 1012; aldicarb, phorate and BAY 25141 controlled root-lesion nematodes on leatherleaf fern, 1016
---: DBCP (1,2-dibromo-3-chloropropane) inhibited growth of three Pythium spp.

and two Phytophthoras in culture, and in the field inhibited Pythium, 622; control of Paratylenchus hamatus under laboratory conditions, 1085
---: DD-MENCS: preplant treatment increased growth of apple trees and reduced nematode populations, 792; DD-MENCS at 35 gal. per acre gave maximum yields of strawberries in West Virginia, 923;
---: DD-MENCS, 1,3-D, and MBR-CP-PBR controlled root-knot nematode in peach seedlings as well as crown gall, Chenopodium and nutsedge, 1014
---: 1,3-D (1,3-dichloropropene): the 1,3-dichloropropene component of the dichloropropene-dichloropropane mixture is about 50 times more toxic than 1,2-dichloropropane to three types of nematodes, 196; 1,3-D and DBCP gave 100% control of Paratylenchus hamatus under laboratory conditions, 1085
---: D-1410 (S-methyl 1-(dimethylcarbamoyl) -N-[(methylcarbamoyl)oxyl]thioformimidate)(DuPont's 1410 nematicide-insecticide), foliar sprays were effective in reduction of Meloidogyne incognita on tobacco, sugar pumpkins, tomato, sweetpotato, and with Pratylenchus scribneri on pole beans for periods ranging from 21 to 28 days, 187
---: methyl bromide: suppressed cereal root eelworm (Heterodera avenae) in wheat and increased yield, in Greece, 134; fumigation of pine nursery beds controlled Pratylenchus penetrans, 572
---: several tested together: controlled Pratylenchus penetrans on leatherleaf fern transplant material, either as bare-root dip, soil drench, or foliar spray, 287; reduced populations of Meloidogyne hapla on lettuce on organic soil in New York, 368; to control root-knot nematodes in Caladium tubers, 411; soil treatment with pesticides controlled Pratylenchus penetrans but increased severity of weather fleck of tobacco, 672; to control Heterodera cacti on Zygocactus truncatus, 692; to kill eggs in cysts of Heterodera tabacum, 781; as chemical dips to control Meloidogyne incognita on dogwood seedlings, 952; for control of root-knot nematode on gardenia, 966
Nematodes: improved method for concentrating nematodes from the Baermann apparatus, 899; Anguina tritici in combination with Corynebacterium tritici and Heterodera tritici found to cause wheat diseases in India, 594
---: citrus-root, citrus nematodes from around olive roots were more infective to olive than biotype -1, or -2 collected from around citrus roots, 717
---: Criconemoides rusticum parasitic on shortleaf pine roots in Georgia, 465
---: cyst (see also golden): cactus cyst, Heterodera cacti controlled on Zygocactus truncatus with nematicide drenches, 692
---: ---: cereal root eelworm (Heterodera avenae), methyl bromide reduced population in wheat, in Greece, 134
---: ---: soybean, Heterodera glycines, new races found capable of attacking formerly resistant varieties, 319; four races known to occur in the United States are characterized, 544
---: ---: sugarbeet, found in cabbage field in Florida, and apparently the first occurrence in the southeastern United States, 635
---: ---: tobacco, rate of increase of Heterodera tabacum in soil is affected by level of population of Pratylenchus penetrans, 25; six non-volatile nematicides effective against emerged larvae, but did not kill many eggs inside cysts, 781
---: dagger: Xiphinema bakeri associated with a serious root disease (corky root) of Douglas-fir seedlings, 165
---: Ditylenchus dipsaci and Pratylenchus penetrans cultured monoxenically, 251
---: golden: Heterodera rostochiensis was not found in Connecticut potato fields in summer of 1969, 261; Rindite, a material used for breaking the rest period of potatoes gave 98 to 100% control of encysted golden nematode larvae in New York test, 306
---: pin: Paratylenchus hamatus from California failed to increase on celery or mint, previously reported hosts in other parts of the world, 920; P. hamatus reported from Iran, 1085
---: plant-parasitic, associated with citrus decline in southern Iran, 339
---: reniform, Rotylenchulus reniformis reported for the first time from Lebanon on tomato and bean, 435
---: root-knot: Meloidogyne incognita acrita found pathogenic to 28 varieties of kenaf, but not to roselle, in Nigerian tests, 547; on lettuce, M. hapla reduced in muck soils of New York with nematicides, 368; chemical treatments with four nematicides found to be as effective as a hot water treatment of 50°C for 30 minutes for control of Meloidogyne spp. in Caladium tubers, 411; root-knot and lesion, M. incognita and Pratylenchus controlled on several host plants with foliar spray or sprays of DuPont 1410,

187; root-knot, sugarbeet, and lesion, the 1,3-dichloropropene components of dichloropropene-dichloropropane mixtures primarily responsible for toxicity to nematodes, 196; M. naasi a parasite of turfgrass in California, 940; M. incognita controlled on gardenia with nematicides, 966; pathogenicity of M. naasi to 'Wintok' oats established, 972; control on peach seedlings with DD-MENCS, 1,3-D, and MBR-CP-PBR, 1014; root-knot and reniform, completely controlled on sweetpotato rootlets after hot water treatment, 1056
---: root lesion: control of Pratylenchus penetrans on leatherleaf fern with several nematicides, 287; P. penetrans, controlled in pine nursery with methyl bromide, 572; P. penetrans, control with preplant treatments increased apple tree growth, 792; P. penetrans, granular applications of phosphates and carbamates for control on Easter lily, 1012; low populations of both P. penetrans and Tylenchorhynchus claytoni found in Connecticut potato fields in 1969, 995; P. penetrans controlled on leatherleaf fern with nematicides, 1016
---: several together: Hoplolaimus galeatus, Helicotylenchus pumulis, Tylenchorhynchus cylindricus, Xiphinema americanum, and X. diversicaudatum parasitized Pinus ponderosa seedlings under greenhouse conditions for 13 months, 752
Neovossia indica: 391
Nepal: 1092
New diseases: undetermined disease of filbert (Corylus avellana), a potential threat to the filbert industry in Oregon, 69; needle cast on bristlecone fir, for the first time in California, 275; stem rot of Ageratum, new in the United States, 1095
New distribution: bacterial blight, of soybean found in New Zealand, 86
---: bean common blight and fuscous blight, occurred in New Zealand in 1970, traced to use of imported seed, 1068
---: bunch disease of pecan, found for first time in Georgia, 12
---: Ceratocystis species (4) not previously reported from Minnesota caused cankers on aspen, 460
---: Cercospora spot on tobacco in United Arab Republic, 666
---: of Dutch elm disease, in Texas, 931
---: Gloeosporium rot of citrus found in Florida, 173
---: Helminthosporium spiciferum found in Sudan for the first time infecting Vicia faba plants, 405
---: liver spot disease of pecan found in Alabama, 408
---: Macrophomina phaseoli, cause of wilt of sunflower, in Hungary, 286
---: maize dwarf mosaic reported in Arizona for first time, 179
---: mistletoe: limber pine dwarf mistletoe found on Pinus flexilis in Montana, 441
---: needle cast, Swiss needle cast, recently discovered in Michigan and Wisconsin, 612
---: nematodes: Paratylenchus hamatus found in Iran, 1085
---: ---: Diphtherophora communis reported from Iran, in citrus orchards, 339
---: ---: Rotylenchulus reniformis found in Lebanon on tomato and bean in 1969, 435
---: ---: sugarbeet cyst nematode in Florida, 635
---: pecan canker, reported for the first time from Arizona, 179
---: Pelargonium rust in Florida, 827
---: potato diseases, reported for the first time from Peru include black leg, brown rot, pink rot, Macrophoma blight, Virus S, purple top, and cork (potato mop-top virus), 315
---: powdery mildew, of peach found in Virginia during the 1968 growing season, 686
---: root and stem rot of roselle, caused by Phytophthora parasitica, in Nigeria, 23
---: Scleroderris lagerbergii, found on red and jack pine in Minnesota, 132
---: Sclerotinia laxa isolated from grape, apparently a first record of this pathogen in New Zealand, 504; benomyl-tolerant strain of Sclerotinia sclerotiorum on muskmelon in Israel, 909
---: stem pitting of peach, from Illinois, 67; from Greece, 73
---: stem rot of soybean, in Arizona, 179
---: Sugarbeet root rot, in Arizona, 179
---: white pine blister rust and limber pine dwarf mistletoe on limber pine in Montana, 441
New Hampshire: 360
New hosts: of alfalfa mosaic virus in Argentina, Ocimum basilicum and Pitraeae cuneato ovata, 722
---: Alternaria alternata caused leaf spot on mung bean, 453
---: anthracnose of oriental poppy, 1037
---: brinjal mosaic disease, Trianthema monogyna and Boerhaavia diffusa discovered as weed hosts of a new strain of cucumber mosaic 1, 437
---: bromegrass mosaic virus, previously unreported, 191
---: Chalaropsis thielavioides pathogenic to

Poinsettia, 451
---: Clitocybe root rot on pecan, in Georgia, 620
---: Corynebacterium michiganense pathogenic to sweet pepper under field conditions in Israel, 804
---: Cylindrocladium crotalariae reported to be pathogenic to yellow poplar, 320
---: Fomes annosus found on dead Rhododendron maximum and Kalmia latifolia in western North Carolina, in October, 1968, 560
---: Pseudomonas syringae isolated from Austrian winter pea, 467
---: Pyrenochaeta terrestris, a root pathogen of creeping bentgrass, 506
---: sugarcane mosaic virus, on bamboo, Arundinaria gigantea, 815
---: Verticillium albo-atrum caused wilting of mango in Florida, 569
---: Verticillium dahliae, cause of wilt of eggplant in Turkey, 241
---: of watermelon mosaic virus, Trichosanthes dioica found to be natural host, 727
New Jersey: 1037
New South Wales (see Australia)
New York: 101; 105; 306; 368; 395; 744; 771; 783; 792; 853
New Zealand: 86; 504; 1068
Nigeria: 23; 520
North Carolina: 119; 125; 183; 846
Nutrition: high nitrogen levels on 'Merion' Kentucky bluegrass and low moisture levels increased Fusarium roseum infection, 1018

Oak: live oak decline, Cephalosporium spp. is major component of disease complex, Hypoxylon atropunctatum accounts for another portion, and Dothiorella quercina may be part of life cycle of Cephalosporium wilt in Texas, 669; virus-like disorder of overcup oak (Quercus lyrata) seedlings from acorns collected in Alabama, no causal agent found, 844; vascular mycosis, cause of oak wilt in eastern Europe, 905
Oats: field studies of injury due to air pollution, 8; stem rust, Puccinia graminis f. sp. avenae, race 6AF comprised 74% of total isolates tested for determination of the physiological races in the United States in 1968, 256; leaf blotch, Pyrenophora avenae shows tolerance to organic mercurial seed treatments, 783; sensitivity of crown rust fungus to insecticide, parathion, 807; reaction of 47 varieties of oats and species of Avena to Meloidogyne naasi in monoxenic culture, 972
Ocimum basilicum (basil): found to be natural host of alfalfa mosaic virus in Argentina, 722
Ohio: 8; 156; 240
Oil: summer oil emulsion used to test effect of spray on spread of strawberry viruses in Oregon, 479
Olive: Infectivity of three biotypes of citrus nematode on two varieties of olive, 'Manzanillo' more tolerant than 'Ascolano', 717
Onion: onion and alfalfa callus produced on Krusberg's and White's media for monoxenic culture of plant parasitic nematodes, 251; onion smut, effect of planting depth and thiram formulations on stands and incidence of smut in field-seeded onions, 290
Ophiostoma (Ceratocystis) roboris: 905
Opuntia tuna (cactus): witches'-broom disease, electron microscopy revealed evidence of existence of three viruses and mycoplasma-like bodies in diseased Opuntia tuna, 851
Oregon: 69; 479; 488; 651; 700
Orosius argentatus: 903
Ozone: high oxidant levels in Cincinnati, Ohio caused plant injury in the field to several plant species, 8; ozone and sulfur dioxide synergism, injury to ponderosa pine geographic race plantation in Puget Sound region, 945; 35 accessions or cultivars of Phaseolus vulgaris tested for tolerance to ozone injury, 1026

Pacific Northwest: 701
Pakistan: 630
Pansy: Pythium myriotylum inoculum from irrigation pond pathogenic to pansy, 1077
Papaya (Carica papaya): distortion ringspot virus, induced apocarpy and "double papaya" fruit formation in several papaya cultivars, in India, 181; susceptibility of cultivars to papaya ringspot and papaya mosaic viruses, 893
Paratylenchus hamatus: 920; 1085
Parsley (Petroselinum crispum): western celery mosaic virus found in parsley crops in Britain, 370; severe stunting and yellowing of plants in Britain due to nasturtium ringspot virus, 734
Pea: Austrian winter pea, a new host of Pseudomonas syringae, 467
---: damping-off: seedling loss due to Pythium sp. and Rhizoctonia solani reduced with fungicidal treatments, 349; caused by Rhizoctonia solani, seed dressing at sowing or soil drenching considered best treatments for control, 1049
---: virus: fizzletop disease, host effects and diagnostic symptoms characterized

by plant dwarfing, leaf twisting, uneven ovule development, 355; a seedborne virus detected in plant samples and seed lots in Washington and Idaho was not found in fresh processing peas in 1969 from seed produced in the Pacific Northwest, 497
Peach (see also Prunus): bacterial spot: foliage injury by dodine and dodine-captan sprays, captan alone is not toxic but appears to be the toxic agent in the spray combination, 152; Xanthomonas pruni controlled with weekly sprays of antibiotic, oxytetracycline, in Maryland, 208; relative resistance of 26 peach cultivars to bacterial spot and Valsa canker, 395
---: nematode control in nursery stock with various nematicides, 1014
---: peach brown rot, evaluation of fungicides for control in New South Wales, 776
---: peach canker, Cytospora spp. predominant over other suspect pathogens, in Illinois, 832
---: postharvest decay, of fruits as affected by fungicide dips and dip temperatures, 764
---: powdery mildew of peach in Virginia, 686
---: stem pitting: reported from Illinois in trees purchased from an Eastern nursery, 67; reported from Greece, 73
Peanut (Arachis hypogaea): Cercospora leaf spot, benomyl effective for control, 955
---: collar rot, caused by Aspergillus niger, of 20 previously tested varieties, only one, U-4-47-7 (EC 21115) from Sudan was significantly resistant in Indian test, 337
---: pod breakdown, Pythium myriotylum more effective than Rhizoctonia solani as pathogen under most conditions, 840
---: rust, from 173 accessions screened for resistance to two rust cultures in greenhouse test, PI 314817 and PI 315608 were physiologically resistant, 381
---: stunt virus, apparently overwinters in Virginia in white clover; isolating peanut fields appears to reduce disease incidence, 935
Pear: blister canker, cultures of Helminthosporium papulosum f. pyri, inhibited most by maneb, ferbam and thiram, 296
---: fire blight: three inoculation techniques to test for fire blight resistance evaluated, 96; scoring system for rating severity in large populations of trees, 835
---: leaf chlorosis of 'Magness' and 'Bartlett' pears, 871
---: postharvest decay: in 'D'Anjou' pears controlled effectively with preharvest spray of benomyl, 647; control of Penicillium expansum with postharvest dips of benomyl and thiabendazole, Alternaria tenuis not controlled, 655; of Anjou pears reduced with fungicide dips of benomyl or thiabendazole, 887
---: stony pit virus, found in older (30 or more years) plantings in eastern New York, 101
Pecan: bunch disease, found in Mitchell County, Georgia, 12; Clitocybe root rot reported on pecan for the first time in the fall of 1969, in Georgia, 620; Cytospora canker, disease characterized by cankers on scaffold and other branches, recorded for the first time in Arizona, possibly in the U.S., 179; downy spot, benomyl, dodine and triphenyltin hydroxide gave best control and also controlled scab, 725; liver spot disease, found in several counties in Alabama, 408; scab, inoculation studies with pecan scab fungus, Fusicladium effusum, 829
Pelargonium hortorum: Pelargonium rust found for first time in Florida, 827
Pencillium sp.: 232
--- digitatum: 325
--- expansum: 12; 647; 655; 887
Peniophora gigantea: application of P. gigantea oidia to cut pine stumps for biological control of Fomes annosus, 799
Pennsylvania: 277; 1080
Pepper: Corynebacterium michiganense found to cause fruit and leaf spot disease, 804; Capsicum frutescens var. conoides 'Tabasco', host of tobacco etch virus in Venezuela, 344; several strains of tobacco etch virus found in California, 786
Peridermium harknessii: 853
Peronospora farinosa: 762
--- manshurica: 901
Peru: 315
Petunia: air pollution damage in Cincinnati related to high oxidant levels in atmosphere, 8
Pezicula alba: 640
--- corticola: 640
--- malicorticis: 640
Phaeocryptopus gäumanni: 612
Phaseolus aconitifolius (see Moth bean or Bean)
Phaseolus aureus (see Mungbean or Bean)
Phaseolus vulgaris (see Bean)
Philippines: 516; 565; 1007; 1045
Phleospora robiniae: 939
Phomopsis sp.: 422
Phoradendron bolleanum subsp. pauciflorum: 15
--- juniperinum subsp. libocedri: 15
--- villosum subsp. villosum: 15
Phyllactinia corylea: 399
Phyllody: of safflower, considered to be

a mycoplasma disease, 735
Phyllosticta sp.: 277; 281
--- fraxinicola: 939
--- robiniae: 939
Phymatotrichum omnivorum: 324
Physiogenic disease: leaf necrosis of spinach, prolonged wet cloudy conditions appear to be principal factor for infection, 155
Physiological breakdown: of cranberry, intermittent warming during cold storage inhibits breakdown, 219
Physiological diseases: virus-like disorder of overcup oak possibly of genetic origin or a physiologically induced disease, 844
Phytophthora sp.: 261
--- spp.: 622
--- citricola: 583
--- citrophthora: 608
--- erythroseptica: 315
--- fragariae: 667; 739; occurrence of new race, A-10 in California, 969
--- infestans: 681; 768
--- nicotianae var. parasitica: 608
--- palmivora: 1035
--- parasitica: 23
Phytotoxicity: of ozone to bean cultivars and accessions, 1026
Picea breweriana (Brewer spruce): sugar pine dwarf mistletoe and hemlock dwarf mistletoe pathogenic to rare tree in Oregon and California, 488
Pink rot: of potato, caused by Phytophthora erythroseptica in Peru, 315
Pinus banksiana (jack pine): first report of Scleroderris canker in Minnesota, 132
--- echinata (shortleaf pine): nematode, Criconemoides rusticum parasitic on pine roots in Georgia, 465
--- flexilis (limber pine): white pine blister rust and limber pine dwarf mistletoe discoveries extend the distribution range of these diseases in Montana, 441
--- jeffreyi (Jeffrey pine): Peridermium harknessi identified in Michigan and New York from pine-to-pine inoculations and germ tube characteristics, 853
--- nigra (Austrian pine): effective control of Dothistroma needle blight with Bordeaux, TC-90, and benomyl + polyvinyl chloride, in Iowa tests, 679
--- ponderosa (western yellow pine): Fomes anonsus detection in stump samples incubated on wire racks, saved labor and time over method of wrapping discs in moist paper, 438; fumigation with methyl bromide in nursery beds infested with root-lesion nematodes resulted in more and larger seedlings, 572; five of seven nematode species tested reproduced on P. ponderosa, 752; variation in chlorotic spotting of needles of 10 geographic races thought to be caused by synergistic effects involving both ozone and sulfur dioxide, 945
--- radiata (Monterey pine): pine seedlings grafted with scions from 5-, 10-, 15-, and 25-year-old pines and inoculated with Dothistroma pini, showed more resistance to infection with increasing age of the scion-donating-trees, 403
--- resinosa (red pine): Scleroderris canker infection in Minnesota, 132; control of Scleroderris canker by fungicide sprays of maneb or ziram, 663; outbreak of Lophodermium pinastri in the Lake States caused serious damage to forest nursery stock, 731
--- strobus (eastern white pine): sodium and calcium contamination from run-off or airborne particles from winter salting of highway, 1021; chemical control of white pine blister rust with use of carriers, 1031
--- sylvestris (Scotch pine): 14 forest nurseries suffered severe damage from the needlecast fungus, Lophodermium pinastri, maneb fungicide used for control, 731
--- taeda (loblolly pine): direct stump treatment with Peniophora gigantea established P. gigantea to the exclusion of Fomes annosus, 799
Piricularia oryzae (Pyricularia oryzae): 520; 632; presence on rice seed in the United States, 931; pathogenic races derived from monoconidial cultures, 1045
Pitraea cuneato ovata: a weed in Argentina found to be a natural host of alfalfa mosaic virus, 722
Plant Introductions: of corn, some resistance to Diplodia stalk rot found in 42 introductions, 624
Planting depth: increased depth of planting onion seed increased incidence of smut, however, thiram seed treatment in addition to insecticide + thiram gave control at 1.5-inch planting depth, 290
Plasmodiophora brassicae: 453; 456
Plastic tubing: for draining wetwood trees, in Alabama, 66
Pleospora sp.: 700
Pod breakdown: of peanut, Rhizoctonia solani versus Pythium myriotylum as pathogens, 840
Podosphaera leucotricha: 81
Poinsettia: Thielaviopsis root rot, good control with benomyl, 328; Chalaropsis thielavioides found to be weak pathogen on poinsettias, 451
Poland: 81; 640
Polyporus zonalis: 322

Poppy (Papaver orientale): anthracnose, caused by Gloeosporium sp., 1037
Populus spp.: direct application of cycloheximide to Hypoxylon cankers on aspen arrested activity of cankers for 5 years, 462; four new species of Ceratocystis found on aspen in Minnesota, not previously reported, also Cenangium singulare, 460
Postharvest decay: of apple, fruit decay from Nectria galligena reduced by benomyl in Ireland when used as postharvest dip or preharvest spray, 83
---: of banana fruit, controlled with benomyl and thiabendazole under commercial conditions, in Samoa and New Zealand, 93
---: of peaches, effects of fungicide dip treatments and dip temperatures on postharvest decay, 764
--: of pear: preharvest sprays of benomyl on 'D'Anjou' pears superior to ziram in control of postharvest decay, 647; Penicillium expansum greatly reduced in stored pears with 10-second dip of benomyl or thiabendazole, Alternaria tenuis not controlled, 655; Penicillium rot of Anjou pears controlled with fungicide dips of benomyl or thiabendazole, 887
---: of strawberries, modified atmosphere, chemical, and heat treatments to control decay of California strawberry fruit, 431
Potato: diseases in Peru, 1965-1968, 315; late blight, evaporation rate as an indicator for potato late blight development in plots of different foliage density, 768; soft rot, caused by Erwinia carotovora, ethylenediaminetetraacetic acid (EDTA) decreased soft rot of potato plugs inoculated with E. carotovora, 863
---: nematodes: golden nematode, none found in Connecticut potato fields in summer of 1969, 261; Rindite (chemical to break the rest period) found to control encysted golden nematode larvae, 306; survey of 28 potato fields in Connecticut showed low population of meadow and tobacco stunt nematodes in 1969, 995
---: Verticillium wilt: foliar symptoms greatly reduced by soil drench of benomyl plus Tween 20 in Connecticut tests, 171; infection reduced and tuber yield increased from seed-piece treatment with benomyl, 254
Powdery mildew: 134 new records of powdery mildews on previously unreported host species, collected principally from the California Bay area, 399
---: of apple, caused by Podosphaera leucotricha, systemic action of benomyl protected unsprayed as well as sprayed leaves of apple, 81
---: of cowpea (Vigna sinensis), benomyl, as a soil drench, seed treatment or foliar spray, controlled Erysiphe polygoni, 346
---: of cucurbits: none of 18 cucumber varieties tested were resistant to infection caused by Erysiphe cichoracearum, 52; effect of benomyl and certain cucumber viruses on growth, powdery mildew, and element accumulation by cucumber plants in greenhouse, 141; of muskmelon, field control by root application of benomyl, bioassay method established correlation between disease incidence and the amount of benomyl in plants, 232; of watermelon fruits in South Carolina, 372; of muskmelons, control with benomyl soil drench, 909
---: of hydrangea, benomyl controlled both powdery mildew and Botrytis blight, 606
---: of peach, causal species could not accurately be determined, 686
---: of rose seedlings, cycloheximide and O-199 (as a dip) effectively eradicated powdery mildew, 598
---: of strawberry, caused by Sphaerotheca maculata, 883
---: of wheat, controlled by soil-applied benomyl, in Canada, 91
Pratylenchus penetrans: 25; 196; 251; 287; 572; 672; 792; 995; 1012; 1016
Preharvest spray: control of postharvest decay of 'D'Anjou' pears with benomyl, 647
Preservation: cultures of Botryosphaeria ribis and Dibotryon morbosum were stored under sterile mineral oil at 2-5°C for 8 and 5 years respectively, without loss of viability and B. ribis found pathogenic to 'Golden Delicious' apples, 217
Protein content: infection of Johnson grass and sorghum-Sudan grass hybrid with maize dwarf mosaic virus had little effect on protein content, 424
Pruning: to control Dutch elm disease, unsuccessful in Michigan, 985
Prunus spp. (see also common names): bacterial canker, of French prune, incidence significantly higher on southerly trunk exposures than on all other exposures, also pruning wounds can be potential sites of entry for pathogen, 332
---: stem pitting of peach, found in Illinois, 67; in Greece, 73
---: cherry leaf spot, controlled with benomyl, 81
---: virus: eleven of 21 Prunus test plants other than sweet cherry, found to be hosts of cherry short stem virus, 651; necrotic ring spot virus transmitted

through almond (Prunus amygdalus) seed, 822; systemic invasion of Prunus budsticks by the green ring mottle virus, 917; 12 Prunus viruses tested, only Prunus ringspot virus caused branch- or top-killing of Kwanzan flowering cherry, 1073
Pseudomonas sp: 263
--- cichorii: 964
--- fluorescens: 863
--- glycinea: 86
--- marginalis: 964
--- marginata: 289
--- phaseolicola: 939
--- solanacearum: 315; 516
--- syringae: 332; 467; 939
--- tabaci: 713
Pseudoperonospora cubensis: 52; 108
Pseudotsuga menziesii: nematode populations of Xiphinema bakeri significantly larger in areas of British Columbia forest nurseries where a serious root disease (corky root) of Douglas-fir occurred, 165; Swiss needle cast reported from the Lake States for the first time, 612
Psychrometer: an electrical psychrometer that senses and records relative humidity at different heights, in widely spaced locations and in confined places, also a computer program for system is available, 19
Puccinia arachidis: 381
--- carthami: 364
--- coronata avenae: 807
--- graminis f. sp. avenae: 256
--- --- f. sp. tritici: 256; 391; 959
--- pelargonii-zonalis: first report from Florida, 827
--- penniseti: 646
--- recondita: 391
--- sorghi: 788
--- striiformis: 347; 384; 391; 992
Puerto Rico: 212
Pyrenochaeta terrestris: 506
Pyrenophora avenae (Helminthosporium avenae): tolerance to organic mercurials, 783
Pyricularia oryzae (see Piricularia)
Pythium sp.: 261; 349; 1077
--- spp.: 8 species isolated and identified from necrotic roots of strawberries in Illinois plantings, 49; 268; 419; 622
--- aphanidermatum: 179
--- myriotylum: 840; 1077
--- splendens: 300
--- sylvaticum: 416
--- ultimum: 1010

Quarantine: quarantine laboratory designed to prevent escape of imported plant pathogens during infectivity tests on experimental crop hosts, 3

Races: 10 geographic races of ponderosa pine show differences with respect to air pollution damage, 945
---: of Aphanomyces euteiches, degree of root damage depends upon bean fungus interaction, 943
---: of Helminthosporium maydis, cause of southern corn leaf blight, 708; physiologic races T and O and disease resistance, 819; 1099; 1100; 1103; 1109; 1114; 1118; past occurrence and geographical distribution of Race T, 1123; 1127
---: new race of Phytophthora fragariae, incitant of red stele of strawberry found in California and designated race A-10, 969
---: race O, of Phytophthora infestans, test for single gene resistance to race O in tomato, 681
---: Puccinia graminis f. sp. tritici, race 15B-2 accounted for 52% of isolates identified from uredial collections in 1968 in the United States, 256
---: Puccinia striiformis, system for differentiating races in the United States, 992
---: of Pyricularia oryzae, identification of races in Nigeria, 520; monoconidial subcultures differentiated into many pathogenic races based on reaction of Philippine and International differentials, 1045
---: soybean cyst nematode: during the growing season of 1969 cyst nematode strains that develop quite freely on roots of resistant varieties were found on farms in Missouri and Tennessee, 319; of soybean cyst nematode in the United States, 544
---: of soybean mildew fungus, sources of immunity to nine races of the fungus, Peronospora manshurica, 901
---: new race of Xanthomonas cyamopsidis found in Texas that is virulent to certain field-resistant varieties of guar, 111
Range: known range of four Ceratocystis species attacking quaking aspen extended to Minnesota, 460
Raspberry: virus diseases of red raspberry, ringspot disease, raspberry mosaic and calico disease found in the Pacific Northwest, 701
Rate: rate of upward movement of Ceratocystis ulmi in elms, 58
Red stele: of strawberry, macerated mycelium from 9 races of fungus (Phytophthora fragariae) were infective to strawberry roots with inoculum dilution of

100:1, 667; inoculation of seedlings with pathogen, P. fragariae, 739
Red thread: of turfgrass, caused by Corticium fuciforme, controlled with benzimidazoles, 1080
Relative humidity: an electric psychrometer for remote sensing and recording of relative humidity, 19
Resistance: in bean: screening for resistance to Rhizoctonia solani in snap bean, 40; in bean, to Fusarium solani f. phaseoli, varieties with purple hypocotyls were more resistant to fungus than those lacking the color; possible relationship of phenolic compounds to resistance, 698; of corn introductions to Diplodia stalk rot, 624; in cucumber, varietal resistance to three cucumber foliar diseases in Florida, 52; of kenaf, to Meloidogyne incognita, none seen in 28 varieties tested, 547; of lettuce to corky root rot, 'Oak Leaf' highly resistant, 'Marquette', a head lettuce type, has high tolerance, 754; of peanut, greenhouse screening for resistance to peanut rust, Puccinia arachidis, 381; in pear seedlings, to fire blight, inoculation techniques tested for screening progeny, 96; in rice varieties, to grassy stunt disease, 565; in roselle, 8 varieties of roselle (Hibiscus sabdariffa) resistant to Meloidogyne incognita acrita, 547; in tomato, to tomato anthracnose, PI 272636 accession from Costa Rica shows some promise as a source for resistance, 32; screening for single gene resistance in tomatoes to race O of Phytophthora infestans, 681
Rhizoctonia spp.: 268
--- solani: 40; 76; 246; 349; 524; 679; 840; 981; 1049; 1080
Rhizoglyphus echinopus: 289
Rhizopus sp.: 647
--- stolonifer: 431; 486; 764
Rhode Island: 168
Rhododendron: indole butyric acid (hormone powder) mixed with benomyl or thiabendazole increased rooting of cuttings significantly, 427; rosebay rhododendron susceptible to Fomes annosus in western North Carolina, apparently a new host, 560
Rice: blast disease, caused by Pyricularia oryzae, 11 races identified in Nigeria, 520; nitrogen fertilization and the rice blast disease, 632; the pathogen, P. oryzae found on 8 of 25 samples examined from the Southern U.S. in 1968 and 1969, 931; great variability in pathogenicity of P. oryzae shown by derivation of races from monoconidial cultures, 1045
---: brown leaf spot disease, caused by Cercospora oryzae, occurs on leaves of Panicum repens in India and may help perpetuate and spread the disease, 272
---: grassy stunt disease, screening method for testing resistance, 565
Roadside trees: tolerance to sodium chloride varies greatly within as well as between tree species, appears to be related to amount of chloride accumulated by leaves of the tree, 360
Root necrosis (see Necrosis)
Root rot (see Rot)
Rose: surface drench of DuPont's 1410 nematicide-insecticide to established greenhouse roses controlled lesion nematode, Pratylenchus scribneri, 187; powdery mildew, caused by Sphaerotheca pannosa var. rosae controlled with cycloheximide and O-199, when used as dips, 598; yield increase of heat-treated 'Baccara' roses in a commercial greenhouse, 825; Rosa setigera, infected with tobacco streak virus in Wisconsin, a woody host, 949
Roselle (Hibiscus sabdariffa): root and stem rot, Phytophthora parasitica was primary cause of infection in Nigeria, 23; eight varieties of roselle tested in Nigeria showed resistance to the root-knot nematode, 547
Rot: bacterial soft rot of lettuce, caused by Erwinia carotovora and Pseudomonas sp., is serious market disease of head lettuce, 263
---: basal rot of bulbous iris, treating with benomyl or thiabendazole within 1 to 2 days rather than 7 days improved control, 235; Fusarium basal rot of Narcissus, benomyl and thiabendazole increased yields of healthy bulbs as well or better than mercurial fungicides, 377
---: boll rot of cotton: cotton boll black rot, Diplodia gossypina controlled with fungicides mixed with an insecticide, 29; estimate of loss from cotton boll rot, 324; internal fluorescence of cotton plants from Arizona found to be unrelated to fiber fluorescence associated with Aspergillus flavus boll rot, 408; microorganisms associated with cotton boll rots in Georgia, 586; mode of entry of boll rot pathogens in cotton varieties and strains, 719; relative frequency of different fungi associated with boll rot, 796
---: brown stem rot of soybean, caused by Cephalosporium gregatum, 987
---: charcoal rot of soybean, incited by Macrophomina phaseoli, found to be transmitted by seeds, 1088
---: charcoal stem rot of sunflower, 891

Rot: citrus brown rot, restricted occurrence in Florida is apparently related to limited distribution of Phytophthora citrophthora and infrequency of climatic conditions causing long periods of fruit wetness, 608
---: collar, of peanut, caused by Aspergillus niger, one variety from Sudan found to be significantly resistant to the disease in comparison to other varieties tested in India, 337
---: corky root rot of lettuce, caused by toxic materials from decomposing lettuce residues; Marquette, a selection with high tolerance has been released, 754
---: corm rot of gladiolus: caused by Fusarium oxysporum f. sp. gladioli, corm treatment with benomyl more effective than thiram, 289; benomyl and thiabendazole fungicide dips reduced number of diseased plants, 689
---: crown of turfgrass, caused by Helminthosporium sp., controlled with benzimidazoles, 1080
---: ear rot of corn, corn ear inoculation with ear rot fungi, Gibberella zeae and Diplodia maydis, 658
--- fruit: of giant granadilla (Passiflora quadrangularis), caused by Botryodiplodia theobromae, 809
---: ---: of strawberry: caused by Botrytis cinerea, reduced in Washington trials with captan, 132; postharvest decay spoilage by B. cinerea and Rhizopus stolonifer reduced most with refrigeration at 35° or exposure to high CO_2 atmosphere for 36 hours, 431; caused by B. cinerea, 883
---: Gloeosporium, of 'Temple' oranges, in Florida, 173
---: gray mold, of lettuce, caused by Botrytis cinerea, ranks with bacterial soft rot in importance as disease in western lettuce on the New York produce market, 263
---: peach brown rot, caused by Monilinia fructicola, captan, dithianon and ziram gave best control in New South Wales, 776
---: red rot of sugarcane: caused by Colletotrichum falcatum, its incidence and spread on midribs of sugarcane leaves in India, 226; disease caused heavy loss in Burma, 427
---: root: of bean, plant material soil amendments and fungicides benomyl and DMTT suppressed Thielaviopsis basicola infections, 114; varieties with purple hypocotyls and black seed showed more resistance to Fusarium solani f. phaseoli than those lacking this color, 698
---: ---: of black walnut, caused by Phytophthora citricola, 583
---: ---: of pea, differential bean series effective in detecting physiological variants of Aphanomyces euteiches, 943
---: ---: of pecan, caused by Clitocybe tabescens, first report on pecan, 620
---: ---: Phymatotrichum omnivorum, loss estimate of cotton crop for 1969, 324
---: ---: of poinsettia, Chalaropsis thielavioides minor component of root rot complex, 451; Thielaviopsis root rot, soil drenches applied to newly-potted rooted cuttings and established plants with root rot symptoms reduced rot effectively, 328
---: ---: of safflower: caused by Pythium splendens, susceptibility of varieties affected by soil temperature, 300; by P. ultimum, 1010
---: ---: of soybean, caused by Thielaviopsis basicola, in Michigan, 849
---: ---: of strawberry, caused by Pythium sylvaticum, 416
---: ---: of sugarbeet, caused by Pythium aphanidermatum, not previously reported from Arizona, 179
---: ---: of tobacco, Thielaviopsis basicola effectively controlled by drenches of benomyl to potted tobacco plants, 136
---: root and stem, of roselle, caused by Phytophthora parasitica, 23
---: soft rot of potatoes, EDTA, an inhibitor of bacterial macerating enzymes reduced soft rot caused by Erwinia carotovora and Pseudomonas fluorescens, 863
---: sour rot of lemon, caused by Geotrichum candidum, 881
---: stalk: 42 plant introductions of corn among 900 tested, showed some resistance to Diplodia stalk rot, 624; influence of root injury, plant population and soil moisture on development of Cephalosporium stalk rot of corn, 989
---: stem: of Ageratum, caused by Alternaria zinniae, new disease in the United States, 1095
---: ---: of carnations, biological control of Fusarium roseum f. sp. dianthi with Bacillus subtilis, which acts as a biological control agent, 446
---: ---: of chrysanthemum, caused by Rhizoctonia solani, 679
---: ---: of soybean, caused by Sclerotinia sclerotiorum, reported for first time from Arizona, 179
---: stem-end: of banana, caused by Gloeosporium musarum, 93; of Chinese gooseberry (kiwi fruit) shipped from New Zealand, Phomopsis sp. was re-

covered from infected fruit, 422
Rotylenchulus reniformis: 435; 1056
Rust: of wheat, current situation for rusts and smuts of wheat in India, 391
Rust: blister: Comandra, infection and mortality on shortleaf pine in Arkansas forests was higher in direct-seeded areas than in plantations, 471
---: ---: white pine: of limber pine in Montana, 441; evaluation of carriers for chemotherapeutant treatment of blister rust cankers in eastern white pine, 1031
Rust: coffee: caused by Hemileia vastatrix, found in Brazil, 355; presence in Brazil confirmed, 539
---: crown, of oats, Puccinia coronata avenae sensitive to insecticide, parathion, in greenhouse fumigation, 807
Rust: fusiform: Cronartium fusiforme, a remote sensing and recording system for relative humidity used successfully in Mississippi with fusiform rust research, 19; new sampler described for estimating spore numbers in the air, with collection at specified time intervals and in discrete zones, 498
---: gall; western gall rust present in Jeffrey pine in Michigan and New York, 853
---: peanut, Puccinia arachidis, 173 accessions screened for resistance, 2 found to be physiologically resistant, 381
---: P. carthami, of safflower seedlings, stand and yield of susceptible and resistant lines, 364
---: P. pelargonii-zonalis observed for first time on geranium in Florida, 827
---: P. penniseti, rapid formation of telial sori in some isolates and none in others could indicate a genetically controlled character in P. penniseti, 646
---: stem: of wheat, annual survey of physiological races of P. graminis f. sp. tritici in the United States in 1968, 256; relationship of spore production of pathogen with temperatures above 21°C coupled with the presence of dew, 959
---: stripe: of wheat, fall infection by stripe rust induced consistent reductions in the number of tillers produced by wheat plants in the Pacific Northwest, 347; natural infection during the summer in Washington occurred when 3 or more hours of rain or dew were recorded, indicating presence of inoculum, 384; wheat plants grown under water stress combined with infection with stripe rust reduced root growth more severely than either factor alone, 387

Safflower (Carthamus tinctorius): Pythium root rot: apparently soil temperature affects the relative susceptibility of safflower varieties to the fungus, 300; susceptibility to Pythium root rot appeared related to period of elongation of hypocotyl-first-internode and not to age alone, 1010; rust, Puccinia carthami, yields in resistant lines were not significantly reduced, 364; safflower phyllody, a mycoplasma disease, in Israel, 735
Salt contamination: of white pine, planted adjacent to an interstate highway, 1021
Salt injury: 22 species of roadside trees examined for sodium chloride tolerance from winter salting of roads in New Hampshire, 12 could be classified as salt-tolerant, and 10 as salt intolerant, 360
Samoa: 93
Sample size: requirements for sample size in testing seed protectants DCNA + captan and DCNA + Difolatan on 15 crop plants, in Michigan, 1041
Scab: of apple, ascospore discharge of Venturia inaequalis on old leaves controlled with benomyl in early spring, 27; scab first found in the Yakima Valley of Washington in 1963, has not been found since, 63; bacterial scab of gladiolus, partially controlled with corm treatments of benomyl and thiram, 289; of pecan, Success cultivar appears to be more susceptible than Stuart cultivar to infection in inoculation studies, 829
Scleroderris lagerbergii: 132; 663
Sclerospora sorghi: 311; 603
Sclerotinia homoeocarpa: 981; 1080
--- laxa: 504
--- sclerotiorum: 28; 179; tolerant strain, 909
--- trifoliorum: 811
Sclerotium rolfsii: 223
Scolytus multistriatus: 266
Scoring system: for severity of fire blight in pear, 835
Scurf: of sweetpotato, hot water treatment at 50°C for 3 to 5 minutes eliminated root-knot and reniform nematodes and reduced scurf on 'Centennial' sweetpotato plants, 1056
Seed: Helminthosporium spiciferum infection of Bermuda grass causes black fungal growth and failure of germination, 358; Pyricularia oryzae occurred or rice seed, particularly from Louisiana, 931; seed protectant field trials, sampe size requirements, 1041; seed corn for 1971, required and recommended labeling procedures to designate the cytoplasm type, that is, nor-

mal, blended or Texas male-sterile, 1114; seed corn production methods in relation to Race T of Helminthosporium maydis, differences in line susceptibility, availability of normal-cytoplasm seed, and the danger of loss from use of F_2 seed or "crib" seed, 1118
---: transmission: of cucumber mosaic virus by infected chickweed plants, 150; of necrotic ring spot virus in almond, 822
---: treatments: roller type mill designed to treat small lots of seeds with liquid or dust formulations, or to mix liquids, inoculum, or chemicals uniformly in small batches of soil, 414; potassium azide, an herbicide, shows promise as a seed treatment, 604; for Pyrenophora avenae on oats, apparent increase in tolerance to organo mercurials observed in New York, 783; for Ustilago nuda, two experimental seed treatments eliminated smut from barley cultivar, Larker, 927
Seedborne: seed infection of chickpea (Cicer arietinum) by Ascochyta rabiei in Israel, 442; a seedborne virus in peas grown in Washington and Idaho, 497; virus-infected seeds in alfalfa seed lots from 4 western States, tests indicate infected seed is cause of initial infection in alfalfa stands, 591; Macrophomina phaseoli found to be carried on and in seed coat of soybean, 1088
Seedling: seedling reaction of inbreds and single crosses of maize to Fusarium moniliforme, 307; corn seedlings inoculated with a toxin extracted from F. moniliforme, showed retarded emergence and growth, 483; use of rice seedlings for testing resistance to grassy stunt disease, 565; Pinus ponderosa seedlings had greater survival and larger plants after bed fumigation with methyl bromide, 572; infection of sorghum seedlings with Sclerospora sorghi by placing infected corn and sorghum leaves in direct contact with seedlings, 603; nematicidal dips controlled Meloidogyne incognita on dogwood seedlings, 952
Seedling diseases: black walnut root rot, caused by Phytophthora citricola, 583
---: of cabbage, systemic infection of 7-day-old cabbage plants by Plasmodiophora brassicae common under greenhouse conditions, 453
---: damping off: of clover, infection by Fusarium, Pythium, Rhizoctonia spp. and a non-sporulator was greatly reduced after 5-minute treatment with radio-frequency energy, 268
---: ---: of cotton: damping-off of cotton seedlings, controlled with benomyl, 76; loss for 1969, 324; controlled with 1,4-oxathiin derivative, 524
---: of pea, reaction of Pythium sp. and five isolates of Rhizoctonia solani to 22 fungicide formulations, 349
---: of safflower: seedling rust, caused by Puccinia carthami, 364; effect of seedling age on Pythium root rot of safflower, 1010
---: of yellow birch seedlings and associated pathogens, 16
Sensitivity: of Paratylenchus hamatus to nematicides in Iranian laboratory experiments, 1085
Septoria sp.: 315
--- apiicola: 531
--- nodorum: 594
--- tritici: 594
Serology: of soybean mosaic virus SMV-M, 301
Setaria viridis: varieties Robust White and Robust Purple differentiate strains A and B of maize dwarf mosaic virus, 630
Smut: brown loose smut (Ustilago nuda), on barley cultivar, Larker, 927
---: loose smut of wheat in India, 391
---: onion, on field-seeded onions, thiram formulations with insecticide + seed treatment with thiram used at three planting depths controlled Urocystis colchici, 290
---: stripe: of 'Merion' Kentucky bluegrass, any treatment sequence in experiments which included 6 oz or more of benomyl 50% WP in either October or March during dormant season resulted in almost complete suppression of smut symptom development, 146; of 'Merion' Kentucky bluegrass, fall application of benomyl gave best protection, but was ineffective against Helminthosporium leaf spot, 168
Snapdragon (Antirrhinum majus): Pythium myriotylum from irrigation pond pathogenic to snapdragon trap plants, 1077
Snow mold: Typhula blight of bentgrass controlled with chloroneb, 1075
Sogata orizicola (Sogatodes orizicola): 56
Soil: amendments: alfalfa hay, cabbage stems and leaves, corn stover, chestnut tannin, lecithin, and oat straw, oilseed meals and plant oils reduced root rot of bean, caused by Thielaviopsis basicola, 114; comparison of systemic activity of thiabendazole and benomyl against Rhizoctonia solani and Sclerotinia homoeocarpa in the greenhouse, 981
---: assay, for isolation of Phytophthora and Pythium species from naturally infested soils in Iran, 261
---: treatments: for fungi: benomyl reduced

symptom development of Verticillium wilt in potato plants in Connecticut, 171; radio-frequency energy to control seedling damping-off in greenhouse soils, 268; benomyl as soil drench controlled powdery mildew of cowpea but treatment caused stunting, 346; benomyl controlled clubroot of cabbage seedlings, 456; Vitavax mixed in cotton field soil controlled damping-off of cotton seedlings from Rhizoctonia solani, 524; to control Fusarium wilt of tomato and watermelon, 561; fungicide applications to the soil, ineffective against powdery mildew of rose seedlings, 598; DBCP has fungicidal properties against several pythiaceous fungi in agar culture and in the field, 622; sand, rather than soil was better for growing strawberry seedlings after inoculation with Phytophthora fragariae, 739; benomyl drench to control powdery mildew of muskmelon, 909; benomyl drench for control of Rhizoctonia damping-off of peas, 1049

---: ---: for nematodes: fumigation with dichloropropene-dichloropropane mixtures, 1,3-dichloropropene component is the part responsible for toxicity to root-knot, sugarbeet, and lesion nematodes, 196; experimental nematicide Dupont 1410 as soil drench effective against Pratylenchus penetrans on leatherleaf fern transplants in Florida, 287; fumigation for controlling root knot of lettuce on organic soil, 368; control of root-lesion nematodes in Pinus ponderosa seedlings with methyl bromide, 572; benomyl reduced nematodes and weather fleck, four nematicides reduced nematodes but increased fleck, 672; soil drenches with several nematicides controlled Heterodera cacti on Christmas cactus, 692; six non-volatile nematicides ineffective against cysted eggs of Heterodera tabacum, but in treated soil, root invasion was curbed and good control attained, 781; around apple trees, in New York, 792; nematicidal drenches gave excellent control of Meloidogyne incognita around gardenia plants, 966; DD-MENCS, 1,3-D, and MBR-CP-PBR effective against Meloidogyne incognita on peach seedlings, 1014; granular applications and soil spray with nematicides controlled root-lesion nematodes on leatherleaf fern, 1016

---: ---: soil fumigation plus fungicidal sprays increased strawberry yield, 923

Solanum melongena (see Eggplant)

Sorghum: Colletotrichum graminicola on Sorghum bicolor introductions, 779 screened for resistance, 60; infection of 'Combine Kafir-60' by Claviceps purpurea after inoculation with the fungus, 844

---: downy mildew: Sclerospora sorghi has spread rapidly in the United States and is now found in 10 States, 311; conidia used as inoculum to inoculate sorghum seedlings with S. sorghi, 603

---: maize dwarf mosaic virus (MDMV): grain sorghum lines and hybrids found to react differently to strains A and B of the virus, 627; reaction of sweet sorghum varieties to MDMV at two locations in Kentucky, 723; severity of MDMV development in Arizona at high elevation and cool summer temperature, 1064

South Africa: 1003

South Carolina: 372

South Dakota: 191

Southern corn leaf blight: an apparent relationship of male-sterile cytoplasm in corn and susceptibility to Helminthosporium maydis, 501; corn seedlings with T or P cytoplasm for male sterility were susceptible to H. maydis, seedlings with normal cytoplasm of S or C were resistant, 708; physiologic races of H. maydis, 819; introduction of the southern corn leaf blight session at the annual meeting of the American Phytopathological Society, 1099; history of the disease, 1100; hosts, 1103; origin and spread in 1970, 1104; physiological races and disease resistance, 1109; toxicity of infected corn or silage to livestock, 1111; protective fungicides, 1112; labeling of 1971 seed, 1114; corn seed production methods in relation to Race T of pathogen, and future seed prospects, 1118; introduction of papers on epidemiology and forecasting, 1122; past occurrence and geographical distribution of isolates of Race T, 1123; perithecia of Cochliobolus heterostrophus on corn leaves in Florida (the perfect stage of H. maydis), 1127; preliminary investigation of blight epiphytotic of 1970, 1129; epidemiology of southern corn leaf blight, exploratory experiments, 1131; plans for forecasting corn blight epidemics, 1134

Southern pea (see Cowpea)

Soybean: bacterial blight, caused by Pseudomonas glycinea, in New Zealand, 86

---: brown stem rot, distribution and isolation of Cephalosporium gregatum in Georgia, 987

---: charcoal rot disease transmitted by seeds, 1088

---: downy mildew resistance, PI 171443, PI 201422, and 'Pine Dell Perfection' immune to nine races of the fungus, 901

---: cyst nematode, physiological strains of Heterodera glycines capable of development on resistant varieties are present in Arkansas, Missouri, and Tennessee, 319
---: root rot, caused by Thielaviopsis basicola, 849
---: stem rot, found in Arizona, caused by Sclerotinia sclerotiorum, 179
---: virus: soybean mosaic virus isolated from cultivar Hood was identified as strain SMV-M and was found to be related to a type strain SMV-NC, 301; cowpea mosaic virus, vector efficiencies of bean leaf beetle and cucumber spotted beetle reduced on soybean test plants, 1053
Sphaerotheca fuliginea: 141; 372; 399
--- humuli: 399
--- lanestris: 399
--- macularis: 362
--- maculata: 883
--- pannosa var. rosae: 598
Spinach: leaf necrosis, atypical climatic conditions of heavy rainfall and humidity thought to be possible cause of disorder found in Texas, 155
Spore dispersal: of Venturia inaequalis ascospores, reduced by fungicidal sprays in early spring on overwintered leaves, 27; new continuous spore sampler designed to collect spores on surface in discrete zones at time intervals, 498
Sporulation: of Pseudoperonospora cubensis, the causal organism for downy mildew of watermelon, 108
Spruce (see Picea)
Statice (Limonium sinuatum): disease of statice in south Florida, 759; control of Cercospora leafspot with several fungicides, 760
Stem breakage: of hackberry in Alabama, caused by Polyporus zonalis, 322
Stem dieback: of yellow birch seedlings, 16
Stemphylium spp.: 580
Stem pitting: of peach, found in 5-year-old peach orchard in Illinois, 67; of peach, in Greece, symptoms closely resemble disorder in the U.S., 73
Storage: physiological breakdown in cranberries greater when held constantly in cold than when shifted to 70° for one day out of each 4 weeks' storage, 219; DCNA-dip effectively reduced occurrence of Rhizopus rot of sweetpotatoes ('Yellow Jersey') with storage temperatures above 59°F, 486; of Stemphylium species under sterile mineral oil, this method may be useful for long-term preservation of Stemphylium species, 580; storage decay of 'D'Anjou' pears, 647
Strains: strain of soybean mosaic virus identified as SMV-M by studies on host range, symptomatology, physical properties, serology, electron microscopy, inclusions and cross protection, 301; cucumber mosaic virus 1, a mosaic disease of brinjal (eggplant) in India found to be new strain of CMV-1, 437; Verticillium albo-atrum, effect of inoculation technique and strains A, B and 2-C of V. albo-atrum on symptom expression in strawberry, 682; of tobacco etch virus, on pepper, 786; watermelon mosaic virus-2 found pathogenic to entire collection of USDA plant introductions of squash and watermelon, 880; powdery mildew resistant strain of Sclerotinia sclerotiorum observed in Israel for first time, 909
---: maize dwarf mosaic virus: in grain sorghum lines, strain A produced systemic infection and B caused only necrotic local lesions, 627; A and B of MDMV, two varieties of Setaria viridis reported as differential hosts, 630
Strawberry: powdery mildew, Sphaerotheca macularis controlled with foliar sprays of benomyl, 362; postharvest decay of fruit, 36-hour period of refrigeration at 35°F or in high CO_2 atmosphere at 59° reduced losses from decay, 431; symptom expression of X Fragaria ananassa as affected by inoculation technique and strains of Verticillium albo-atrum, 682; combination of nematicidal and fungicidal treatments significantly increased yield in West Virginia, 923
---: Botrytis cinerea: captan fungicide increased marketable yields an average of 1500 pounds and net profits $125 per acre, in Washington, 132; Botrytis fruit rot and powdery mildew, fungicidal control, 883
---: Phytophthora fragariae: macerated mycelium of P. fragariae effective for inoculation of strawberry plants, nine races tested were infective at 100:1 dilution, 667; for P. fragariae inoculation, growing seedlings in sand after a mycelial dip, 739; two pure cultures of P. fragariae isolated in California are designated race A-10 of the fungus because of the ability to infect two clones of Fragaria chiloensis, 969
---: Pythium sylvaticum: most common of 8 species of Pythium isolated from diseased strawberry roots in Illinois, 49; P. sylvaticum found pathogenic on strawberry roots, 416
---: virus: virus-free Dabreak and Headliner cultivars compared with compar-

able virus-infected plants, data indicated Headliner was susceptible to mottle virus, Dabreak was tolerant but was susceptible to virus complex found in commercial plantings in Louisiana, 293; relation of oil sprays to the incidence of crinkle, mild yellow edge, necrotic shock, and mottle viruses in field-grown plants in Oregon, 479
Stripe smut: of Kentucky bluegrass, inflorescences infected with Ustilago striiformis, 206
Stump infection: by Fomes annosus, a method for detection of infection in conifer stumps, 438
Sudan: 405; 528
Sugarbeet: root rot, Pythium aphanidermatum pathogenic to first commercial plantings of sugarbeets in Arizona, 179
Sugarcane: red rot disease: caused by Colletotrichum falcatum, the course of development on plants of different varieties and ages is related to weather, 226; reached epiphytotic proportions in Burma in 1969 on predominant sugarcane variety, Co 419, 427
---: sugarcane mosaic virus, in Louisiana, a new strain I produced the most severe effects on germination and yield of three varieties, strain H was more severe than strain A, 554
Sulfur dioxide: four cultivars of highbush blueberry showed tolerance to exposure to sulfur dioxide with variability among cultivars, 704; possible ozone-sulfur dioxide interaction responsible for chlorotic spotting of ponderosa pine foliage near La Grande, Washington, 945
Summer biology: of Puccinia striiformis in southeastern Washington, 384
Summer death: of beans, in New South Wales, Australia, 61 bean varieties and breeding lines tested for resistance to leafhopper-transmitted disease, 903
Sunflower: Macrophomina phaseoli, found for the first time in Hungary, sudden wilting after seed pollination is suggested as the primary plant stress, 286; charcoal stem rot, caused M. phaseoli, response to natural infection in Maryland, 891
Surfactants: comparison of various surfactants as adjuvants for the fungicidal action of benomyl on Verticillium albo-atrum, 741
Survey: of diseases affecting yellow birch seedlings in Upper Michigan, 16; physiological races of Puccinia graminis in the U.S. in 1968, 256; for golden nematode, in Connecticut in 1969, 261; potato disease survey in Peru for 1965-1968, 315; nematode survey of citrus orchards in Iran, 339; for bark canker and phloem canker of Persian walnut in California, 373; Comandra blister rust incidence in Ozark and Ouachita National Forests in Arkansas, 471
Survival (see also Longevity): long-term preservation of Botryosphaeria ribis and Dibotryon morbosum under sterile mineral oil, 217
Sweetpotato: Rhizopus rot, DCNA dip effective in reduction of rot during post storage when holding temperatures were 60°F or above, 486; a suscept of false broomrape, identity of etiological agent not known, 979
---: nematodes: effective control of Meloidogyne incognita with DuPont 1410 as foliar spray, but not as a dip, 187; elimination of root-knot and reniform nematodes from rootlets and reduction of scurf by hot water treatment, 1056
Sweet william (Dianthus barbatus): used as trap plant for Pythium inoculum from irrigation ponds, 1077
Symptomatology: gross symptomatology of Kentucky bluegrass inflorescences infected by Ustilago striiformis, 206; of soybean mosaic virus SMV-M, 301; of Pseudomonas syringae on Austrian winter pea, 467; of Corynebacterium michiganense on sweet pepper, 804
Symptom progression: of Dutch elm disease in inoculated trees, 127
Systemic distribution: of fungicides in sprayed 'Jonathan' apple trees, 996

Taiwan: 29
Target spot: of cucumber, no varieties in screening test found resistant to Corynespora cassiicola, 52
Telia production: identification of different strains of Puccinia penniseti on basis of ability to produce telia appears to be promising, 646
Temperature: effect of dip temperatures of water or fungicides to protect peach fruit from Monilinia fructicola and Rhizopus stolonifer infection, 764; thermotherapy successful in complete cure of whole plant from grapevine fanleaf virus, in France, 915
---: air: relation of temperature with toxicity of dodine-captan sprays on peach, 152; optimum growth of Cercospora sp. on Ficus elastica occurred at 25°C with daily alternation from 10 to 30°, 199; cool summer temperatures and absence of susceptible varieties are major factors limiting geographical distribution of bark and phloem cankers of Persian walnut in California, 373; hot air treatment of virus-tested 'Bac-

cara' roses increased yield over commercial stock that was not heat-treated or virus-tested, 825; development of stem rust spore production favored by temperature above 21° C with minimum of 8 hours of dew, 959; little leaf of apple seedlings, evidence suggests a genetic disorder caused by high midsummer temperature, in Maryland, 1059; low air temperatures related to serious disease incidence of MDMV in sorghum hybrids in Arizona at high elevations, 1064

---: hot water treatment: nematicide treatments compared with hot water for control of root-knot nematodes in Caladium tubers, 411; eleminated root-knot and reniform nematodes and greatly reduced scurf on sweetpotato rootlets, 1056

---: soil: greenhouse soil exposed to radiofrequency energy (rf) for 5 minutes resulted in rise in temperature of soil and subsequent control of damping-off fungi, 268; effect of soil temperature on Pythium root rot of safflower, 300; optimum temperature for reproduction of Meloidogyne naasi was 27°C, 940; high soil temperature and high nitrogen levels correlated with severity of infection by Fusarium roseum, of 'Merion' Kentucky bluegrass, 1018

---: storage: cranberries held at 33 or 38°F and warmed to 70° one day for every 4 weeks' storage had less physiological breakdown than under constant cold storage, 219; postharvest decay of strawberries, refrigeration at 35°F for 36 hours was effective in reducing decay, hot, moist air treatment at 106° followed by cooling at 32° reduced decay but fruit injury was evident, 431; 60°F or above and a DCNA-dip preserved keeping qualities of 'Yellow Jersey' sweetpotatoes during the post storage period, 486

Tennessee: 319

Terminology: terminology and identity of infraspecific forms of the soybean cyst nematode, 544

Tetranychus urticae: 1029

Texas: 108; 111; 155; 311; 931

Thielaviopsis basicola: 114; 136; 223; 328; 849

Tillering: tiller reduction in wheat as result of stripe rust infection in the Pacific Northwest, 347

Tilletia caries: 391

--- foetida: 391

Tobacco: increase in growth of shade tobacco around new cedar posts, cause not determined, 536

Tobacco: air pollution: tobacco types and varieties varied in their reaction to ozone injury, 8; weather fleck, several nematicides increased incidence of fleck but benomyl gave good control until late in season, 672

---: Cercospora spot, found on tobacco leaves in United Arab Republic, apparently first report from this country, 666

---: nematodes: control of Meloidogyne incognita with foliar sprays of DuPont 1410, 187

---: root rot, benomyl superior to 10 other fungicides in control of Thielaviopsis basicola, 136

---: virus: tobacco etch virus: results indicate that some currently grown varieties of flue-cured tobacco in North Carolina differ in their reaction to TEV, 119; host of TEV in Venezuela, 344

Tomato: upper-surface bronzing of leaves, caused by oxidants in air, 8

---: anthracnose, caused by Colletotrichum coccodes, resistance in tomato introductions, 32

---: bacterial canker: stem inoculations of 14-day-old plants with Corynebacterium michiganense gave rapid and uniform disease development, 36; difference in virulence of C. michiganense isolates, 976

---: Fusarium wilt: symptoms reduced by benomyl and correlation with a bioassay of fungitoxicant in tomato plants, 12; controlled in Greece with soil drench of benomyl, 561

---: graywall, tobacco mosaic virus possible cause, 678

---: late blight, detached leaf method for screening tomato plants and at the same time preserving all plants and permitting true replication of a series of tests, 681

---: nematodes: Meloidogyne incognita controlled with experimental nematicide, 187; reniform nematode collected from rhizosphere of tomato, in Lebanon, 435

---: Pythium myriotylum from irrigation ponds pathogenic to tomato seedlings, 1077

--- virus: found to be host of tobacco etch virus in Venezuela, 344

Toxicity: of sulfur dioxide to highbush blueberry, 704; of silage of corn infected with Helminthosporium maydis to livestock, 1111

Toxin: toxin produced by Fusarium moniliforme found to be heat stable and completely water soluble, tests indicate that the pathogenicity of the fungus is

dependent upon its ability to produce a toxin that causes seedling blight of corn, 483
Translocation: evaluation of chemicals for systemic activity against two plant pathogens, by use of carrot discs, 223
Transmission: seed transmission of cucumber mosaic virus by chickweed, 150; of pea fizzletop virus, pea-aphid-transmissible, 355; of brinjal mosaic disease, 437; transmissible and non-transmissible apple fruit-blotching syndromes, 490; of cotton leaf curl virus with use of static cage technique, 528; grassy stunt disease of rice, insect vector, Nilaparvata lugens used to inoculate seedlings in mass screening test for resistance in rice varieties, 565; seed transmission of strawberry latent ringspot virus by celery seed, 802; of necrotic ring spot virus through almond seeds, 822; Myzus persicae transmitted maize dwarf mosaic virus from corn to corn with greater frequency than sugarcane mosaic virus when single aphids were used, 876; of rice blast disease, untreated rice seed may be source of primary inoculum in the U.S., 931; of charcoal rot of soybean, by seeds, 1088
--: mechanical: two graft-transmissible bark diseases of apple, pustule canker and freckle scurf, reported from Washington, 44; knife for making leaf-tissue grafts to transmit citrus viruses, 907
Trianthema monogyna: weed host of brinjal mosaic disease, a new strain of cucumber mosaic virus 1, 437
Trichosanthes dioica: found to be natural host of watermelon mosaic virus, in India, 727
Tsuga heterophylla (western hemlock): field tests indicated that use of two herbicides in thinning young stands of western hemlock did not prevent infection by Fomes annosus, 330
Tundu: of wheat, in India, caused by nematode, Anguina tritici and bacterium, Corynebacterium tritici, 594
Turkey: 241; 881
Twig dieback (see Dieback)
Tylenchorhynchus claytoni: 672; 995
--- cylindricus: 752
Tylenchulus semipenetrans: 339; 717
Typhula itoana: 1075

Umbelliferous crops: western celery mosaic virus found in celery, coriander and parsley crops in Britain, 370
United Arab Republic: 666
Urethane foam: additives, durability, and expansion of a urethane foam useful in tree cavity fills, 858
Urocystis colchici: 290
--- tritici: 391
Ustilago nuda: 927
--- striiformis: 146; 168; 206
--- tritici: 391
Utah: 364

Valsa ambiens: 16
--- cincta: 395
--- leucostoma: 395
Variegation disorder: wheat spindle streak mosaic present in virtually all winter wheat grown in Michigan in 1969, 635
Vascular mycosis: of oak, in Europe, 905
Vectors associated with: brinjal mosaic disease, Aphis gossypii and Myzus persicae, 437; cassava mosaic virus, Bemisia tabaci, 34; cowpea mosaic virus, Cerotoma ruficornis transmits virus, 212; retention of infectivity of bean leaf beetle and spotted cucumber beetle, 1053; Dutch elm disease, Scolytus multistriatus, 266; 706; grassy stunt disease of rice, Nilaparvata lugens, 565; greening disease of citrus, Diaphorina citri, 1092; maize dwarf mosaic virus and sugarcane mosaic virus, Myzus persicae, 876; peanut stunt virus, Aphis craccivora, 935; safflower phyllody, leafhopper, Neoaliturus fenestratus, 735; summer death of beans, Orosius argentatus, 903; tobacco etch virus, Aphis craccivora, 344; western celery mosaic virus transmitted by Cavariella pastinacae, C. aegopodii, and Myzus persicae, 370; wheat spot chlorosis, Aceria tulipae is vector of disease resembling wheat spot mosaic virus, in Ohio, known properties still too few to establish relationship, 156; wheat streak mosaic virus, Aceria tulipae, 635
Venezuela: 344
Venturia inaequalis: 27; 63
Vermont: 638
Verticillium albo-atrum: 12; 171; 254; 324; 508; 569; 682; 741; 846
--- dahliae, wilt of eggplant, first report from Turkey of infection on this crop, 241
--- nigrescens: 586
Viability (see Survival)
Vigna sinensis (see Cowpea)
Virginia: 686; 935; 955
Virus: alfalfa mosaic virus: apparently most of initial AMV infections in alfalfa stands are due to infected seed, 591; alfalfa mosaic virus, on basil (Ocimum basilicum) and a weed, Pitraeae cuneato

ovata in Mendoza, Argentina as natural hosts, 722
---: apple: evidence of a relationship between green mottle, chat fruit, and flat apple viruses of apple, 105; two graft-transmissible fruit blotch disorders in British Columbia presumed to be virus diseases, 490; fruit wrinkle, a transmissible disease of 'Newtown' apple in British Columbia thought to have been introduced from eastern Canada in 'Gano' nursery stock imported in 1904, 493
---: of barley, barley stripe mosaic, infection levels and field spread in 'Compana' barley in Montana, 749
---: brome mosaic, spotted spider mite failed to transmit virus from barley to barley, 1029
---: bromegrass mosaic virus, serological, electron microscopic and physical properties of virus are reported as well as host range list including previously unreported hosts, 191
---: cactus (Opuntia tuna), evidence of three viruses and mycoplasma-like bodies associated with witches'-broom disease, 851
---: citrus: tobacco necrosis virus of citrus, by artificial inoculation 389; tristeza virus of citrus in Chile, 542; greening virus disease of citrus in West Pakistan, 630; impietratura disease of citrus, infected budwood grafted to 'Madam Vinous' and 'Sour' orange always yields leaf flecking symptoms on young leaves, this could be valuable as an indexing procedure, 643; use of double-bladed knife for leaf-tissue grafts for transmission of citrus viruses, 907
---: cotton: tobacco ringspot: cotton cv. Coker 201 developed local and systemic symptoms when mechanically inoculated with serological strains of TRSV, 342; transmission of cotton leaf curl virus by single insects of Bemisia tabaci, 528
---: cowpea: cowpea mosaic virus found in Puerto Rico belongs to the beetle-transmitted group of cowpea mosaic viruses, 212; several viruses found internally seedborne in cowpea seed grown for seed, in eight States, 243; Tetranychus urticae failed to transmit tobacco ringspot virus from cowpea or squash to cowpea, 1029
---: cucurbits: cucumber mosaic (CMV), tomato ringspot and sour cherry necrotic ringspot did not influence level of benomyl required to completely suppress powdery mildew on cucumber, 141; CMV transmitted through chickweed seed, 150; CMV isolates from sugarbeet, lettuce and cantaloup showed some variation, 176
watermelon mosaic virus (WMV), Trichosanthes dioica found to be a natural host of this virus in India, 727; epiphytotic of WMV in central New York affecting several species, 744; 30 to 100% of USDA plant introductions of squash (Cucurbita pepo) and watermelon (Citrullus lanatus) were susceptible to WMV-2 in screening test, 880
---: Datura, viruses isolated from 37 arborescent Datura species from Bolivia, Ecuador and Colombia, 675
---: Desmodium mosaic virus, evidence from electron microscopy indicates this virus is a member of the potato virus Y group, 161
---: of gladiolus, survey of North Carolina fields indicates bean yellow mosaic, cucumber mosaic, tomato ringspot virus and tobacco ringspot virus are major viral pathogens of gladiolus in the State, 125
---: grape: grapevine fanleaf, whole plant completely cured of virus by thermotherapy, 915; grape viruses in early California grapevines, 1055
---: maize dwarf mosaic (MDMV): two varieties of Setaria viridis are hosts that differentiate strains A and B of the virus, 630; comparative transmission of MDMV and sugarcane mosaic virus by the green peach aphid, Myzus persicae, 876; on sorghum, 179; 424; 627; 723; 1064
---: nasturtium ringspot virus isolated from parsley crops in Britain, 734
---: papaya: distortion ringspot virus, cause of abnormal formation of fruit, 181; papaya ringspot and papaya mosaic 90 accessions tested were susceptible to both viruses, 893
---: pea: fizzletop disease of pea can be seed-transmitted at 80% frequencies, 355; a seedborne virus of pea detected by indexing in 15 of 53 fresh plant samples having virus-like symptoms and in 4 of 113 seed lots from Washington and Idaho, 497
---: peanut stunt virus, was a very destructive disease to beans in North Carolina in 1969, 183; association of white clover and aphids with the virus, in Virginia, 935
---: pear, stony pit virus of pear, found in six out of seven older plantings in eastern New York, newer plantings were free from the disease, 101
---: pepper, tobacco etch virus on peppers in California, 786
---: of potato, viruses appear to be the most destructive group of potato path-

Virus (continued): ogens in Peru, 315
---: Prunus: cherry short stem virus of sweet cherry, host range studies, 651; necrotic ring spot, passage of virus through almond seeds, 822; green ring mottle virus systemic in budsticks of four Prunus species, 917; top- or branch-killing of Kwanzan flowering cherry by Prunus ringspot virus, 1073
---: red raspberry, ringspot disease, caused by tomato ringspot virus was most common virus in field-grown plants in Oregon and Washington, raspberry mosaic limited mostly to 'Newburgh' in British Columbia, calico disease was rare, 701
---: of rose, heat-treated and virus-tested 'Baccara' roses outyielded commercial stock not treated or virus-tested, 825; tobacco streak virus on Rosa setigera, first report on a woody host, 949
---: sorghum: maize dwarf mosaic: in sorghum and Johnson grass, reported from Arizona for first time, in 1968 and 1969, 179; infection in Johnson grass and a sorghum-Sudan grass hybrid, virus had no detrimental effects on protein content or digestibility by steers, however, it did reduce yield of hybrid for second and third cuttings, not the first, 424; of grain sorghum lines, no cross-protection observed between two strains of the virus, 627; differential reactions of sweet sorghum varieties to MDMV in Kentucky, 723; development in sorghum hybrids in Arizona in relation to low temperature and high elevation, 1064
---: soilborne wheat streak mosaic, inoculation with artist's airbrush significantly increased infection over manual rubbing technique, 550
---: soybean: soybean mosaic virus, SMV-M identified from Glycine max variety Hood, 301; cowpea mosaic virus, vector efficiencies of spotted cucumber beetle and bean leaf beetle reduced on soybean test plants compared with cowpea, 1053
---: of strawberry: effect of virus on fruit yield of Headliner and Dabreak varieties, 293; effect of oil sprays on spread of viruses, in Oregon, 479; strawberry latent ringspot virus transmitted in celery seed, 802
---: sugarcane mosaic: in sugarcane varieties, effects on stands and yields, 554; found on Arundinaria gigantea, a bamboo, in Louisiana, 815
---: tobacco: tobacco etch virus, its effect on yield and quality of flue-cured tobacco varieties, 119; in Venezuela, 344
---: tobacco mosaic virus suggested as a causal agent for graywall of ground tomatoes in Florida, 678
---: western celery mosaic in umbelliferous crops in Britain, 370
---: wheat: soilborne wheat streak mosaic virus, 550; wheat spindle streak mosaic, in Michigan winter wheat, soil transmission of disorder and virus-like particles and pinwheel inclusions suggest a soilborne virus as the causal entity, 635; wheat spot chlorosis pathogen, from Ohio and its relation to wheat spot mosaic virus, 156

Walnut (Persian walnut, Juglans regia): varietal and geographic distribution of two bacterial diseases, bark canker and phloem canker of walnut in California, 373; root rot of black walnut in Indiana nursery, caused by Phytophthora, citricola, 583
Washington: 44; 63; 132; 235; 330; 347; 377; 384; 497; 917; 945; 1073
Watermelon (see Cucurbits)
Water stress: in wheat roots, its influence alone, or when combined with stripe rust in mist culture experiments, 387
Weather: for initiation of midrib lesions on sugarcane caused by Colletotrichum falcatum, wet weather conditions are essential and determine the extent of initiation and spread, 226; epiphytotic of sugarcane by red rot in Burma, abnormally wet weather coupled with possible use of infected seed pieces thought to be cause, 427; relation of weather to development of infection foci on wheat stem rust, 959
Weather fleck: of tobacco, caused by ozone, soil treatment with benomyl decreased severity of fleck, four nematicides tested increased the severity, 672
Weed hosts: for Verticillium albo-atrum, variation in host reaction was found among species of the same family, 508
West Germany: 136
West Virginia: 923
Wetness: measurement of dew periods and their duration with a modified Taylor dew meter, 512
Wetting parameter: use of electrical resistance sensors to record the wetting duration and degree of wetness, 474
Wetwood: of Siberian elm in Alabama: Erwinia nimipressuralis, the common pathogen causes fluxing, plastic tubing is effective as a drain for flux, 66
Wheat: diseases of wheat in India other than rusts and smuts, 594
---: nematodes, infestation by Heterodera avenae reduced with methyl bromide, 134

---: powdery mildew, caused by Erysiphe graminis, benomyl banded into soil with seed gave good control, 91
Wheat: rust: stem: distribution and prevalence of physiologic races of Puccinia graminis in the U.S. in 1968, 256; effects of temperature and dew on size and development of infection foci, 959
---: ---: stripe: fall infection by Puccinia striiformis in the Pacific Northwest resulted in fewer tillers, less straw, and lower yield, 347; the summer biology of Puccinia in southeastern Washington and natural infection during the summer, 384; influence of stripe rust and water stress on wheat roots in mist culture, 387; differential cultivars to identify races of P. striiformis in the U.S., 992
---: virus: wheat spindle streak mosaic, in 1969 practically all Michigan-grown winter wheat was affected, 635; wheat spot chlorosis, also known as "wheat leaf variegation" is similar to wheat spot mosaic virus, but additional properties of pathogen need to be established, 156; wheat streak mosaic virus, 550
Wilt: bacterial, vascular wilt of abaca and banana, caused by Pseudomonas solanacearum, in Philippines, 516
---: Fusarium: of tomato, symptoms reduced by benomyl fungicide, 12; estimated cotton loss from, in 1969, 324; wilt of tomato and watermelon controlled with soil drench of benomyl, 561
---: Macrophomina phaseoli caused wilt of sunflower in Hungary for the first time, pathogen thought to have been introduced into the country from elsewhere, 286
---: oak wilt complex in Europe, disagreement over causal agent or agents, 905
---: Verticillium: of potato, control by soil treatment with benomyl, 171; treatment of potato seed tubers with benomyl reduced wilt and increased yields of tubers, 254; estimated loss of cotton from, 324; reaction of some common cultivars and weeds to an isolate of V. albo-atrum from cotton, 508; of mango, in Florida, apparently a first report of disease in mango, 569; of eggplant, terminal grafts of eggplant on resistant tomato rootstocks yielded high level of resistance in eggplant scion, 846
Wisconsin: 281; 612; 754; 949
Wyoming: 547

Xanthomonas cyamopsidis: 111
--- malvacearum: 324; 713
--- phaseoli: 557; 1068
--- --- var. fuscans: 1068
--- pruni: 208; 395
Xiphinema americanum: 752
--- bakeri: 165
--- diversicaudatum: 752

Yellow poplar (Liriodendron tulipifera): virulence of three Cylindrocladium species on potted seedlings, 320
Yield: virus-free plants of strawberry varieties Dabreak and Headliner compared with fruit yields of virus-infected plants, 293; effect of maize rust (Puccinia sorghi) on different corn varieties and the impact on yield, 788; yield loss of cotton, caused by parasitic diseases in 1969, 324; reduction caused by stripe rust infection of wheat in the fall, 347; reduced yield of safflower, from seedling rust, 364; effect of maize dwarf mosaic virus infection on yield of Johnson grass and a sorghum-Sudan grass hybrid, 424

Zea mays (see Corn)
Zygocactus truncatus (Christmas cactus): nematicides effective in control of cyst nematode, Heterodera cacti, with pot drenches, 692

PLANT SCIENCE RESEARCH DIVISION

Prepared by Marguerite G. Benedict

UNITED STATES DEPARTMENT OF AGRICULTURE
Agricultural Research Service
Beltsville, Maryland 20705

Official Business
Penalty for Private Use, $300

POSTAGE & FEES PAID
United States Department of Agr

EES PAID
ent of Agr